OMEG 2017, International Symposium on Origin of Matter and Evolution of Galaxies, Daejeon, Korea, JUNE 27-30, 2017

Contribution ID: 115

Type: 3 Min. Talk for Poster Presentor

eta-decay scheme of 140 Te to 140 I: Suppression of Gamow-Teller transitions between the neutron $h_{9/2}$ and proton $h_{11/2}$ doublet orbitals

Thursday, 29 June 2017 17:13 (3 minutes)

In astrophysical environments, the beta(β)-decay of unstable nuclei plays a crucial role in generating the final isotopic abundances along with rapid neutron-capture reactions (r-processes). The β -decay strengths are largely influenced by the so-called Gamow-Teller (G-T) interaction that is closely related to the spin-isospin interaction. This spin-isospin excitation is one of important ingredient for modeling the nuclear shell structure. Consequently, the role played by the G-T transition provides plenty of information on nuclear structure and nucleosynthesis. The G-T strength has been mainly constrained by the transformation between a proton and a neutron in the high-angular momentum orbital-spin partners such as; $\pi f_{7/2}$ - $\nu f_{7/2}$, $\pi g_{9/2}$ - $\nu g_{9/2}$, and $\pi h_{11/2}$ - $\nu h_{11/2}$. However, in extreme neutron-rich environments, it can be controlled by the interplay between a neutron in the upper orbital and a proton in the lower orbital among a spin-orbital doublets, for instance, $\pi h_{11/2}$ - $\nu h_{9/2}$.

Investigating of the neutron-rich nuclei beyond ¹³²Sn has attracted much attention because they are expected to reveal changes of size, diffuseness, and shell closures and, on the other side, to give information on nucleosynthesis along the r-process paths. However, it is a challenging task to determine decay rates and life times for the nuclei in this region. Here we present for the first time the β -decay of ¹⁴⁰Te (Z = 52) to odd-odd ¹⁴⁰I (Z = 53). The parent nuclide ¹⁴⁰Te was produced through the in-flight fission of ²³⁸U beam with the energy of 345 MeV per nucleon on a ⁹Be target at the Radioactive Ion Beam Factory (RIBF), RIKEN. By using a β -delayed γ -ray spectroscopy, a decay scheme of ¹⁴⁰Te have been assigned as a 1⁺ state based on the log ft values. These 1⁺ states can be interpreted as being associated with the $\pi h_{11/2} \nu h_{9/2}$ configuration formed by the G-T transition between a neutron in the $h_{9/2}$ orbital and a proton in the $h_{11/2}$ orbital. Systematic features of level structures and G-T transitions are discussed within the framework of the spherical shell model and deformed shell model descriptions.

Presenter: MOON, Byul

Session Classification: Poster session (Chair: K. Cho)