OMEG 2017, International Symposium on Origin of Matter and Evolution of Galaxies, Daejeon, Korea, JUNE 27-30, 2017

Contribution ID: 54

Type: Oral Presentation

Observational constraints on first-star nucleosynthesis

Tuesday, 27 June 2017 09:40 (25 minutes)

Over the past 25 years, it has been recognized that the light element carbon plays a crucial role in the early chemical enrichment of the Universe. One fundamental observation is that the frequency of the so-called carbon-enhanced metal-poor (CEMP) stars in the Milky Way increases dramatically with decreasing iron abundance –from 20% of all stars with [Fe/H] < -2.0 to > 80% of stars with [Fe/H] < -4.0. Recent discoveries of enhanced carbon in damped Lyman alpha systems at high redshift reveal that the abundance patterns observed in this gas are commensurate with a sub-class of the CEMP stars, the so-called CEMP-no stars, which exhibit little or no enhancement of their neutron-capture elements –providing one of the first direct observational linkages between the high-z Universe and presently observed stars in the Galaxy. I summarize recent progress on our understanding of the production of carbon by first-generation stars, and the powerful constraints that this information provides on Galactic chemical evolution models, the initial mass function in the early Universe, and the nature of first-star nucleosynthesis.

Presenter: BEERS, Timothy

Session Classification: Session 1 (Chair: Y. S. Lee)