

# Dark Halo Investigation by DAMA/LIBRA

The 1<sup>st</sup> IBS Conference on Dark World

Daejeon, South Korea

(30<sup>th</sup> October – 3<sup>rd</sup> November 2017)



Vincenzo Caracciolo for the DAMA collaboration.  
National Laboratory of Gran Sasso, INFN.



# DAMA Experimental Activities

**DAMA Collaboration** (spokesperson: **prof. R. Bernabei**):

Roma2, Roma1, LNGS-INFN, IHEP/Beijing

+ by-products and small scale expts.: INR-Kiev and others

+ neutron meas.: ENEA-Frascati e ENEA-Casaccia

+ in some studies on  $\beta\beta$  decays (DST-MAE project): IIT Kharagpur/Ropar, India



DAMA/CRYS  
DAMA/R&D  
DAMA/LXe  
DAMA/Ge

DAMA/NaI  
↓  
DAMA/LIBRA

**DAMA: an observatory for rare processes @LNGS**



# Relic DM Particles from Primordial Universe

## What Accelerators can do:

- to demonstrate the existence of some of the DM candidates

## What Accelerators cannot do:

- to credit that a certain particle is a DM solution or the "only" DM particle solution...

+ DM candidates and scenarios exist (even for neutralino candidate) on which accelerators cannot give any information

- Composition?  
DM multicomponent also in the particle part?

- Right related nuclear and particle physics?

etc... etc...

SUSY  
(as neutralino or sneutrino  
In various scenarios)

the sneutrino in the Smith  
and Weiner scenario

sterile  $\nu$

mirror dark matter

invisible axions,  $\nu$ 's

even a suitable particle not  
yet foreseen by theories

axion-like (light pseudoscalar  
and scalar candidate)

self-interacting dark matter

electron interacting dark matter

Kaluza-Klein particles (LKK)

heavy exotic candidates, as  
"4th family atoms", ...

Elementary Black holes,  
Planckian objects,  
Daemons

etc...

a heavy  $\nu$  of the 4-th family

&

Right halo model and parameters?

Non thermalized components?

Caustics?

clumpiness?




# Some Direct Detection Processes:

- Inelastic Dark Matter:  $W + N \rightarrow W^* + N$ 
  - $W$  has 2 mass states  $\chi_+$ ,  $\chi_-$  with  $\delta$  mass splitting
  - Kinematic constraint for the inelastic scattering of  $\chi_-$  on a nucleus


$$\frac{1}{2} \mu v^2 \geq \delta \Leftrightarrow v \geq v_{thr} = \sqrt{\frac{2\delta}{\mu}}$$

- Elastic scatterings on nuclei
  - detection of nuclear recoil energy




- Excitation of bound electrons in scatterings on nuclei
  - detection of recoil nuclei + e.m. radiation

- **Conversion of particle into e.m. radiation**
  - detection of  $g$ , X-rays,  $e^-$




- Interaction only on atomic electrons
  - detection of e.m. radiation



- Interaction of light DMp (LDM) on  $e^-$  or nucleus with production of a lighter particle
  - detection of electron/nucleus recoil energy

e.g. sterile  $\nu$



e.g. signals from these candidates are completely lost in experiments based on “rejection procedures” of the e.m. component of their rate

... also other ideas ...

- ... and more




# The DM Annual Modulation: a Model Independent Signature to Investigate the DM Particles Component in the Galactic Halo

With the present technology, the annual modulation is the main model independent signature for the DM signal. Although the modulation effect is expected to be relatively small a suitable large-mass, low-radioactive set-up with an efficient control of the running conditions can point out its presence.

## Requirements of the DM annual modulation

- 1) Modulated rate according cosine
- 2) In a definite low energy range
- 3) With a proper period (1 year)
- 4) With proper phase (about June 2<sup>nd</sup>)
- 5) Just for single hit events in a multi-detector set-up
- 6) With modulation amplitude in the region of maximal sensitivity must be <7% for usually adopted halo distributions, but it can be larger in case of some possible scenarios

Drukier, Freese, Spergel PRD86; Freese et al. PRD88



$$v_{\oplus}(t) = v_{\text{sun}} + v_{\text{orb}} \cos \gamma \cos[\omega(t-t_0)]$$

$$S_k[\eta(t)] = \int_{\Delta E_k} \frac{dR}{dE_R} dE_R \cong S_{0,k} + S_{m,k} \cos[\omega(t-t_0)]$$

- $v_{\text{sun}} \sim 232 \text{ km/s}$  (Sun vel in the halo)
- $v_{\text{orb}} = 30 \text{ km/s}$  (Earth vel around the Sun)
- $\gamma = \pi/3$ ,  $\omega = 2\pi/T$ ,  $T = 1 \text{ year}$
- $t_0 = 2^{\text{nd}} \text{ June}$  (when  $v_{\oplus}$  is maximum)

the DM annual modulation signature has a different origin and peculiarities (e.g. the phase) than those effects correlated with the seasons

To mimic this signature, spurious effects and side reactions must not only - obviously - be able to account for the whole observed modulation amplitude, but also to satisfy contemporaneously all the requirements



# The pioneer DAMA/NaI: 100 kg highly radiopure NaI(Tl)

## Performances:

N.Cim.A112(1999)545-575, EPJC18(2000)283,  
Riv.N.Cim.26 n. 1(2003)1-73, IJMPD13(2004)2127

## Results on rare processes:

- Possible Pauli exclusion principle violation
- CNC processes
- Electron stability and non-paulian transitions in Iodine atoms (by L-shell)
- Search for solar axions
- Exotic Matter search
- Search for superdense nuclear matter
- Search for heavy clusters decays

PLB408(1997)439  
PRC60(1999)065501


PLB460(1999)235  
PLB515(2001)6  
EPJdirect C14(2002)1  
EPJA23(2005)7  
EPJA24(2005)51

## Results on DM particles:

- PSD
- Investigation on diurnal effect
- Exotic Dark Matter search
- Annual Modulation Signature

PLB389(1996)757  
N.Cim.A112(1999)1541  
PRL83(1999)4918

PLB424(1998)195, PLB450(1999)448, PRD61(1999)023512,  
PLB480(2000)23, EPJC18(2000)283, PLB509(2001)197, EPJC23(2002)61,  
PRD66(2002)043503, Riv.N.Cim.26 n.1 (2003)1, IJMPD13(2004)2127,  
IJMPA21(2006)1445, EPJC47(2006)263, IJMPA22(2007)3155,  
EPJC53(2008)205, PRD77(2008)023506, MPLA23(2008)2125.



*Data taking completed on July 2002, last data release 2003. Still producing results*


**Model independent evidence of a particle DM component in the galactic halo at  $6.3\sigma$  C.L.**

**total exposure (7 annual cycles) 0.29 ton  $\times$  yr**



# DAMA/LIBRA set-up ~250 kg NaI(Tl) (Large sodium Iodide Bulk for RAre processes)

As a result of a second generation R&D for more radiopure NaI(Tl)  
by exploiting new chemical/physical radiopurification techniques  
(all operations involving crystals and PMTs - including photos - in HP Nitrogen atmosphere)



Residual contaminations  
in the new DAMA/LIBRA  
NaI(Tl) detectors:  
 $^{232}\text{Th}$ ,  $^{238}\text{U}$  and  $^{40}\text{K}$   
at level of  $10^{-12}$  g/g

- **Radiopurity, performances, procedures, etc.:** NIMA592(2008)297, JINST 7 (2012) 03009
- **Results on DM particles: Ann. Mod. Signature:** EPJC56(2008)333, EPJC67(2010)39, EPJC73(2013)2648
- **related results:** PRD84(2011)055014, EPJC72(2012)2064, IJMPA28(2013)1330022, EPJC74(2014)2827, EPJC75 (2015) 239, EPJC75(2015)400, IJMPA 31 (2016) dedicated issue, EPJC77(2017)83
- **Results on rare processes: PEP violation in Na, I:** EPJC62(2009)327, **CNC in I:** EPJC72(2012)1920  
**IPP in  $^{241}\text{Am}$ :** EPJA49(2013)64



# Complete DAMA/LIBRA-phase1

	Period	Mass (kg)	Exposure (kg×day)	$(\alpha - \beta^2)$
DAMA/LIBRA-1	Sept. 9, 2003 - July 21, 2004	232.8	51405	0.562
DAMA/LIBRA-2	July 21, 2004 - Oct. 28, 2005	232.8	52597	0.467
DAMA/LIBRA-3	Oct. 28, 2005 - July 18, 2006	232.8	39445	0.591
DAMA/LIBRA-4	July 19, 2006 - July 17, 2007	232.8	49377	0.541
DAMA/LIBRA-5	July 17, 2007 - Aug. 29, 2008	232.8	66105	0.468
DAMA/LIBRA-6	Nov. 12, 2008 - Sept. 1, 2009	242.5	58768	0.519
DAMA/LIBRA-7	Sep. 1, 2009 - Sept. 8, 2010	242.5	62098	0.515
DAMA/LIBRA-phase1	Sept. 9, 2003 - Sept. 8, 2010		379795 ± 1.04 ton×yr	0.518
DAMA/NaI + DAMA/LIBRA-phase1:			1.33 ton×yr	

- EPJC56(2008)333
- EPJC67(2010)39
- EPJC73(2013)2648
- calibrations:  $\approx 96$  Mevents from sources
- acceptance window eff: 95 Mevents ( $\approx 3.5$  Mevents/keV)

a ton × yr experiment? done

## DAMA/LIBRA-phase1:

- First upgrade on Sept 2008: replacement of some PMTs in HP N<sub>2</sub> atmosphere, new Digitizers (U1063A Acqiris 1GS/s 8-bit High-speed cPCI), new DAQ system with optical read-out installed

## DAMA/LIBRA-phase2 (running):

- Second upgrade at end 2010: replacement of all the PMTs with higher Q.E. ones from dedicated developments
- commissioning on 2011

**Goal: lowering the software energy threshold**

- Fall 2012: new preamplifiers installed + special trigger modules. Other new components in the electronic chain in development






# Model Independent Annual Modulation Result

**DAMA/NaI + DAMA/LIBRA-phase1** Total exposure: **1.33 ton×yr**

EPJC 56(2008)333,  
EPJC 67(2010)39,  
EPJC 73(2013)2648

**Residual rate of the 2-6 keV single-hit scintillation events vs time**



Absence of modulation? No

$$\chi^2/\text{dof} = 154/87$$

$$P(A=0) = 1.3 \times 10^{-5}$$

Fit with all the parameters free:

$$A = (0.0112 \pm 0.0012) \text{ cpd/kg/keV}$$

$$t_0 = (144 \pm 7) \text{ d} - T = (0.998 \pm 0.002) \text{ y}$$

**The data favor the presence of a modulated behaviour with all the proper features for DM particles in the galactic halo at about  $9.3\sigma$  C.L.**



# Model Independent Annual Modulation Result


**DAMA/NaI + DAMA/LIBRA-phase1** Total exposure: 487526 kg×day = **1.33 ton×yr**

EPJC 56(2008)333, EPJC 67(2010)39, EPJC 73(2013)2648


The measured modulation amplitudes (A), period (T) and phase ( $t_0$ ) from the single-hit residual rate vs time

	A(cpd/kg/keV)	T=2 $\pi$ / $\omega$ (yr)	$t_0$ (day)	C.L.
<b>DAMA/NaI+DAMA/LIBRA-phase1</b>				
(2-4) keV	<b>0.0190 ±0.0020</b>	<b>0.996 ±0.002</b>	<b>134 ± 6</b>	<b>9.5<math>\sigma</math></b>
(2-5) keV	<b>0.0140 ±0.0015</b>	<b>0.996 ±0.002</b>	<b>140 ± 6</b>	<b>9.3<math>\sigma</math></b>
(2-6) keV	<b>0.0112 ±0.0012</b>	<b>0.998 ±0.002</b>	<b>144 ± 7</b>	<b>9.3<math>\sigma</math></b>

**Acos[w(t-t<sub>0</sub>)]**



**No systematics** or side reaction able to account for the measured modulation amplitude and to satisfy all the peculiarities of the signature



This result offers an additional strong support for the presence of DM particles in the galactic halo further excluding any side effect either from hardware or from software procedures or from background. The residual rates of the single-hit events measured over the 7 DAMA/LIBRA annual cycles are reported as collected in a single cycle

The data favor the presence of a modulated behaviour with all the proper features for DM particles in the galactic halo at about 9.3 $\sigma$  C.L.



# Model Independent Annual Modulation Result

DAMA/NaI + DAMA/LIBRA-phase1

Total exposure: 487526 kg×day = 1.33 ton×yr

Max-lik analysis of single hit events




EPJC 56(2008)333, EPJC 67(2010)39, EPJC 73(2013)2648

- No modulation above 6 keV
- No modulation in the whole energy spectrum
- No modulation in the 2-6 keV multiple-hit events

$$R(t) = S_0 + S_m \cos[\omega(t - t_0)]$$

here  $T = 2\pi/\omega = 1$  yr and  $t_0 = 152.5$  day

$$R(t) = S_0 + S_m \cos[\omega(t - t_0)] + Z_m \sin[\omega(t - t_0)] = S_0 + Y_m \cos[\omega(t - t^*)]$$




No systematics or side processes able to quantitatively account for the measured modulation amplitude and to simultaneously satisfy all the many peculiarities of the signature are available.



# Rate behaviour above 6 keV

- **No Modulation above 6 keV**



Mod. Ampl. (6-10 keV): cpd/kg/keV

- (0.0016 ± 0.0031) DAMA/LIBRA-1
- (0.0010 ± 0.0034) DAMA/LIBRA-2
- (0.0001 ± 0.0031) DAMA/LIBRA-3
- (0.0006 ± 0.0029) DAMA/LIBRA-4
- (0.0021 ± 0.0026) DAMA/LIBRA-5
- (0.0029 ± 0.0025) DAMA/LIBRA-6
- (0.0023 ± 0.0024) DAMA/LIBRA-7

→ statistically consistent with zero


- **No modulation in the whole energy spectrum:**

- studying integral rate at higher energy,  $R_{90}$
- $R_{90}$  percentage variations with respect to their mean values for single crystal in the DAMA/LIBRA running periods
- Fitting the behaviour with time, adding a term modulated with period and phase as expected for DM particles:

**consistent with zero**

Period	Mod. Ampl.
DAMA/LIBRA-1	-(0.05±0.19) cpd/kg
DAMA/LIBRA-2	-(0.12±0.19) cpd/kg
DAMA/LIBRA-3	-(0.13±0.18) cpd/kg
DAMA/LIBRA-4	(0.15±0.17) cpd/kg
DAMA/LIBRA-5	(0.20±0.18) cpd/kg
DAMA/LIBRA-6	-(0.20±0.16) cpd/kg
DAMA/LIBRA-7	-(0.28±0.18) cpd/kg

DAMA/LIBRA-phase1



$s \approx 1\%$ , fully accounted by statistical considerations

+ if a modulation present in the whole energy spectrum at the level found in the lowest energy region →  $R_{90} \sim \text{tens cpd/kg} \rightarrow \sim 100 \sigma$  far away


No modulation above 6 keV, no modulation in the whole energy spectrum, no modulation in the 2-6 keV multiple-hit events → This accounts for all sources of bckg and is consistent with the studies on the various components



# Summary of the results obtained in the additional investigations of possible systematics or side reactions – DAMA/LIBRA-phase1

(NIMA592(2008)297, EPJC56(2008)333, J. Phys. Conf. ser. 203(2010)012040, arXiv:0912.0660, S.I.F.Attn Conf.103(211), Can. J. Phys. 89 (2011) 11, Phys.Proc.37(2012)1095, EPJC72(2012)2064, arxiv:1210.6199 & 1211.6346, IJMPA28(2013)1330022, EPJC74(2014)3196 )

Source	Main comment	Cautious upper limit (90%C.L.)
RADON	Sealed Cu box in HP Nitrogen atmosphere, 3-level of sealing, etc.	$<2.5 \times 10^{-6}$ cpd/kg/keV
TEMPERATURE	Installation is air conditioned+ detectors in Cu housings directly in contact with multi-ton shield→ huge heat capacity + T continuously recorded	$<10^{-4}$ cpd/kg/keV
NOISE	Effective full noise rejection near threshold	$<10^{-4}$ cpd/kg/keV
ENERGY SCALE	Routine + intrinsic calibrations	$<1-2 \times 10^{-4}$ cpd/kg/keV
EFFICIENCIES	Regularly measured by dedicated calibrations	$<10^{-4}$ cpd/kg/keV
BACKGROUND	No modulation above 6 keV; no modulation in the (2-6) keV <i>multiple-hits</i> events; this limit includes all possible sources of background	$<10^{-4}$ cpd/kg/keV
SIDE REACTIONS	Muon flux variation measured at LNGS	$<3 \times 10^{-5}$ cpd/kg/keV





# No role for $\mu$ in DAMA annual modulation result

## ✓ Direct $\mu$ interaction in DAMA/LIBRA set-up:

DAMA/LIBRA surface  $\approx 0.13 \text{ m}^2$

m flux @ DAMA/LIBRA  $\approx 2.5 \mu/\text{day}$

It cannot mimic the signature: already excluded by  $R_{90}$ , by *multi-hits* analysis + different phase, etc.

## ✓ Rate, $R_n$ , of fast neutrons produced by $\mu$ :

- $\Phi_\mu$  @ LNGS  $\approx 20 \mu \text{ m}^{-2} \text{d}^{-1}$  ( $\pm 1.5\%$  modulated)
- Annual modulation amplitude at low energy due to m modulation:

$$S_m^{(m)} = R_n g e f_{DE} f_{\text{single}} 2\% / (M_{\text{setup}} DE)$$

Moreover, this modulation also induces a variation in other parts of the energy spectrum and in the *multi-hits* events


## ✓ Inconsistency of the phase between DAMA signal and m modulation

$\mu$  flux @ LNGS (MACRO, LVD, BOREXINO)  $\approx 3 \cdot 10^{-4} \text{ m}^{-2} \text{s}^{-1}$ ; modulation amplitude 1.5%; **phase**: July  $7 \pm 6 \text{ d}$ , June  $29 \pm 6 \text{ d}$  (Borexino)

**The DAMA phase: May  $26 \pm 7 \text{ days}$  (stable over 14 years)**


The DAMA phase is  $5.7\sigma$  far from the LVD/BOREXINO phases of muons ( $7.1\sigma$  far from MACRO measured phase)

... many others arguments EPJC72(2012)2064, EPJC74(2014)3196



$$S_m^{(m)} < (0.3-2.4) \times 10^{-5} \text{ cpd/kg/keV}$$

It cannot mimic the signature: already excluded by  $R_{90}$ , by *multi-hits* analysis + different phase, etc.





# No role for n/μ/ν in DAMA annual modulation result

- Contributions to the total neutron flux at LNGS;
- Counting rate in DAMA/LIBRA for *single-hit* events, in the (2 – 6) keV energy region induced by:
  - neutrons,
  - muons,
  - solar neutrinos.

(See e.g. also EPJC 56 (2008) 333, EPJC 72(2012) 2064, JIMPA 28 (2013) 1330022) EPJC74(2014)3196

$$\Phi_k = \Phi_{0,k} (1 + \eta_k \cos \omega (t - t_k))$$

$$R_k = R_{0,k} (1 + \eta_k \cos \omega (t - t_k))$$

Modulation amplitudes

Source	$\Phi_{0,k}^{(n)}$ (neutrons cm <sup>-2</sup> s <sup>-1</sup> )	$\eta_k$	$t_k$	$R_{0,k}$ (cpd/kg/keV)	$A_k = R_{0,k} \eta_k$ (cpd/kg/keV)	$A_k / S_m^{exp}$
SLOW neutrons	thermal n (10 <sup>-2</sup> – 10 <sup>-1</sup> eV)	1.08 × 10 <sup>-6</sup> [15] however ≪ 0.1 [2, 7, 8]	–	< 8 × 10 <sup>-6</sup> [2, 7, 8]	≪ 8 × 10 <sup>-7</sup>	≪ 7 × 10 <sup>-5</sup>
	epithermal n (eV-keV)	2 × 10 <sup>-6</sup> [15] however ≪ 0.1 [2, 7, 8]	–	< 3 × 10 <sup>-3</sup> [2, 7, 8]	≪ 3 × 10 <sup>-4</sup>	≪ 0.03
FAST neutrons	fission, (α, n) → n (1-10 MeV)	≃ 0.9 × 10 <sup>-7</sup> [17] however ≪ 0.1 [2, 7, 8]	–	< 6 × 10 <sup>-4</sup> [2, 7, 8]	≪ 6 × 10 <sup>-5</sup>	≪ 5 × 10 <sup>-3</sup>
	μ → n from rock (> 10 MeV)	≃ 3 × 10 <sup>-9</sup> (see text and ref. [12])	0.0129 [23] end of June [23, 7, 8]	≪ 7 × 10 <sup>-4</sup> (see text and [2, 7, 8])	≪ 9 × 10 <sup>-6</sup>	≪ 8 × 10 <sup>-4</sup>
	μ → n from Pb shield (> 10 MeV)	≃ 6 × 10 <sup>-9</sup> (see footnote 3)	0.0129 [23] end of June [23, 7, 8]	≪ 1.4 × 10 <sup>-3</sup> (see text and footnote 3)	≪ 2 × 10 <sup>-5</sup>	≪ 1.6 × 10 <sup>-3</sup>
	ν → n (few MeV)	≃ 3 × 10 <sup>-10</sup> (see text)	0.03342 * Jan. 4th *	≪ 7 × 10 <sup>-5</sup> (see text)	≪ 2 × 10 <sup>-6</sup>	≪ 2 × 10 <sup>-4</sup>
direct μ	Φ <sub>0</sub> <sup>(μ)</sup> ≃ 20 μ m <sup>-2</sup> d <sup>-1</sup> [20]	0.0129 [23]	end of June [23, 7, 8]	≃ 10 <sup>-7</sup> [2, 7, 8]	≃ 10 <sup>-9</sup>	≃ 10 <sup>-7</sup>
direct ν	Φ <sub>0</sub> <sup>(ν)</sup> ≃ 6 × 10 <sup>10</sup> ν cm <sup>-2</sup> s <sup>-1</sup> [26]	0.03342 *	Jan. 4th *	≃ 10 <sup>-5</sup> [31]	3 × 10 <sup>-7</sup>	3 × 10 <sup>-5</sup>

\* The annual modulation of solar neutrino is due to the different Sun-Earth distance along the year; so the relative modulation amplitude is twice the eccentricity of the Earth orbit and the phase is given by the perihelion.

+ In no case neutrons (of whatever origin), muon or muon induced events, solar ν can mimic the DM annual modulation signature since some of the **peculiar requirements of the signature** would fail (and – in addition - quantitatively negligible amplitude with respect to the measured effect).

All are negligible w.r.t. the annual modulation amplitude observed by DAMA/LIBRA and they cannot contribute to the observed modulation amplitude.

# Final model independent result

## DAMA/NaI+DAMA/LIBRA-phase1

Presence of modulation **over 14 annual cycles at  $9.3\sigma$  C.L.** with the proper distinctive features of the DM signature; all the features satisfied by the data over 14 independent experiments of 1 year each one

The total exposure by former DAMA/NaI and present DAMA/LIBRA is  **$1.33 \text{ ton} \times \text{yr}$**  (14 annual cycles)

In fact, as required by the DM annual modulation signature:

1)

The *single-hit* events show a clear cosine-like modulation, as expected for the DM signal

2)

Measured period is equal to  $(0.998 \pm 0.002) \text{ yr}$ , well compatible with the 1 yr period, as expected for the DM signal

3)

Measured phase  $(144 \pm 7)$  days is well compatible with the roughly about 152.5 days as expected for the DM signal

4)

The modulation is present only in the low energy (2–6) keV energy interval and not in other higher energy regions, consistently with expectation for the DM signal

5)

The modulation is present only in the *single-hit* events, while it is absent in the *multiple-hit* ones as expected for the DM signal

6)

The measured modulation amplitude in NaI(Tl) of the *single-hit* events in the (2–6) keV energy interval is:  $(0.0112 \pm 0.0012) \text{ cpd/kg/keV}$  ( $9.3\sigma$  C.L.).

No systematic or side process able to simultaneously satisfy all the many peculiarities of the signature and to account for the whole measured modulation amplitude is available

# Final model independent result DAMA/NaI+DAMA/LIBRA-phase1

Presence of modulation **over 14 annual cycles at 9.3 $\sigma$  C.L.** with the proper distinctive features of the DM signature; all the features satisfied by the data over 14 independent experiments of 1 year each one

The total exposure by former DAMA/NaI and present DAMA/LIBRA is **1.33 ton  $\times$  yr** (14 annual cycles)

In fact, as required by the DM annual modulation signature:

1)

The *single-hit* events show a clear cosine-like modulation, as expected for the DM signal

2)

Measured period is equal to  $(0.998 \pm 0.002)$  yr, well compatible with the 1 yr period, as expected for the DM signal

3)

Measured phase  $(144 \pm 7)$  days is well compatible with the roughly about 152.5 days as expected for the DM signal

4)

The modul energy (2- in other high expe

5)

The modulation is present only in the *single-hit* events, while it is absent in the *multiple-hit* ones as expected for the DM signal

The meas of the *single-hit* (0.0112

No systematic or side process able to simultaneously the signature and to account for the whole meas

## Model-independent evidence by DAMA/NaI and DAMA/LIBRA

well compatible with several candidates in many astrophysical, nuclear and particle physics scenarios

Neutralino as LSP in various SUSY theories

Various kinds of WIMP candidates with several different kind of interactions  
Pure SI, pure SD, mixed + Migdal effect + channeling, ... (from low to high mass)

a heavy  $\nu$  of the 4-th family

Pseudoscalar, scalar or mixed light bosons with axion-like interactions

WIMP with preferred inelastic scattering

Mirror Dark Matter

Light Dark Matter

Dark Matter (including some scenarios for WIMP) electron-interacting

Sterile neutrino

Self interacting Dark Matter

heavy exotic candidates, as "4th family atoms", ...

Elementary Black holes such as the Daemons

Kaluza Klein particles

... and more





# About Interpretation and Comparisons

See e.g.: Riv.N.Cim.26 ono.1(2003)1,  
IJMPD13(2004)2127, EPJC47(2006)263,  
IJMPA21(2006)1445, EPJC56(2008)333,  
PRD84(2011)055014, JMPA28(2013)1330022

## ...models...

- Which particle?
- Which interaction coupling?
- Which EFT operators contribute?
- Which Form Factors for each target-material?
- Which Spin Factor?
- Which nuclear model framework?
- Which scaling law?
- Which halo model, profile and related parameters?
- Streams?
- ...


## ...and experimental aspects...

- Exposures
- Energy threshold
- Detector response (phe/keV)
- Energy scale and energy resolution
- Calibrations
- Stability of all the operating conditions.
- Selections of detectors and of data.
- Subtraction/rejection procedures and stability in time of all the selected windows and related quantities
- Efficiencies
- Definition of fiducial volume and non-uniformity
- Quenching factors, channeling
- ...

Uncertainty in experimental parameters, as well as necessary assumptions on various related astrophysical, nuclear and particle-physics aspects, affect all the results at various extent, both in terms of exclusion plots and in terms of allowed regions/volumes. Thus comparisons with a fixed set of assumptions and parameters' values are intrinsically strongly uncertain.

**No experiment can - at least in principle - be directly compared in a model independent way with DAMA**

# Is it an “universal” and “correct” way to approach the problem of DM and comparisons?




No, it isn't. This is just a largely arbitrary/partial/incorrect exercise





# ... an example in literature...

## Case of DM particles inducing elastic scatterings on target-nuclei, Spin-Independent case



### Regions in the nucleon cross section vs DM particle mass plane


- Some velocity distributions and uncertainties considered.
- The DAMA regions represent the domain where the likelihood-function values differ more than  $7.5\sigma$  from the null hypothesis (absence of modulation).
- For CoGeNT a fixed value for the Ge quenching factor and a Helm form factor with fixed parameters are assumed.
- The CoGeNT region includes configurations whose likelihood-function values differ more than  $1.64\sigma$  from the null hypothesis (absence of modulation). This corresponds roughly to 90% C.L. far from zero signal.

DAMA allowed regions for the considered scenario without (green), with (blue) channeling, with energy-dependent Quenching Factors (red);

$7.5 \sigma$  C.L.

CoGeNT; qf at fixed assumed value  
 $1.64 \sigma$  C.L.

Compatibility also with first CRESST and CDMS, if the two CDMS-Ge, the three CDMS-Si and the CRESST recoil-like events are interpreted as relic DM interactions



Including the Migdal effect  
→ Towards lower mass/higher  $\sigma$

PRD84(2011)055014,  
IJMPA28(2013)1330022

Co-rotating halo,  
Non thermalized component  
→ Enlarge allowed region towards larger mass

Combining channeling and energy dependence of q.f. (AstrPhys33 (2010) 40) → Towards lower  $\sigma$




# DAMA annual modulation effect and Symmetric mirror matter

Eur. Phys. J. C (2017) 77


Symmetric mirror matter:

- halo composed by a bubble of Mirror particles of different species; Sun is travelling across the bubble which is moving in the Galactic Frame (GF);
- the mirror particles in the bubble have Maxwellian velocity distribution in a frame where the bubble is at rest; cold and hot bubble with temp from  $10^4$  K to  $10^8$  K
- interaction via photon - mirror photon kinetic mixing

Examples of expected phase of the annual modulation signal



The blue regions correspond to directions of the halo velocities in GC ( $\theta, \phi$ ) giving a phase compatible at 3 $\sigma$  with DAMA phase





# DAMA annual modulation effect and Symmetric mirror matter

Eur. Phys. J. C (2017) 77


Symmetric mirror matter:

- Results refers to halo velocities parallel or anti-parallel to the Sun ( $\alpha = 0, \pi$ ). For these configurations the expected phase is June 2
- The only parameter whose value will be varied in the analysis is the  $V_{\text{halo}}$  module (positive velocity will correspond to halo moving in the same direction of the Sun while negative velocity will correspond to opposite direction)

Mirror matter composition	H (%)	He (%)	C (%)	O (%)	Fe (%)
H', He'	25	75	–	–	–
H', He', C', O'	12.5	75.	7.	5.5	–
H', He', C', O', Fe'	20	74	0.9	5.	0.1

DAMA/LIBRA allowed values for  $\sqrt{f}\epsilon$  in different scenarios

where  $f$  is the fraction of DM in the Galaxy in form of mirror atoms and  $\epsilon$  is the coupling constant



Many configurations and halo models favoured by the DAMA annual modulation effect corresponds to couplings values well compatible with cosmological bounds.



# DAMA/LIBRA – phase2

After a period of tests and optimizations in data taking in this new configuration




Second upgrade on Nov/Dec 2010: all PMTs replaced with new ones of higher Q.E.

typically

DAMA/LIBRA-phase1: 5.5-7.5 ph.e./keV

→ DAMA/LIBRA-phase2: 6-10 ph.e./keV




- To study the nature of the particles and features of related astrophysical, nuclear and particle physics aspects, and to investigate second order effects
- Special data taking for other rare processes
- + R&D in progress towards more future phase3




# DAMA/LIBRA – phase2

After a period of tests and optimizations in data taking in this new configuration




more IJMPA28(2013)1330022



Second upgrade on Nov/Dec 2010: all PMTs replaced with new

typically

DAMA/LIBRA-phase1: 5.5-7.5 ph.e./keV  
→ DAMA/LIBRA-phase2: 6-10 ph.e./keV



<sup>234m</sup> Pa (Bq/kg)	<sup>235</sup> U (mBq/kg)	<sup>228</sup> Ra (Bq/kg)	<sup>228</sup> Th (mBq/kg)	<sup>40</sup> K (Bq/kg)	<sup>137</sup> Cs (mBq/kg)	<sup>60</sup> Co (mBq/kg)
-	47	0.12	83	0.54	-	-
-	10	0.02	17	0.16	-	-

- To study the nature of the particles and features of related astrophysical, nuclear and particle physics aspects, and to investigate second order effects
- Special data taking for other rare processes
- + R&D in progress towards more future phase3



# DAMA/LIBRA phase 2 – data taking



Second upgrade at end of 2010: JINST 7(2012)03009  
all PMTs replaced with new ones of higher Q.E.

**Energy resolution** mean value: prev. PMTs 7.5% (0.6% RMS)  
new HQE PMTs 6.7% (0.5% RMS)

- ✓ Fall 2012: new preamplifiers installed + special trigger modules.
- ✓ Calibrations 5 a.c.: ~  $1.03 \times 10^8$  events from sources
- ✓ Acceptance window eff. 5 a.c.: ~  $7 \times 10^7$  events (~  $2.8 \times 10^6$  events/keV)
- ✓ Exposure collected in the first 5 a.c. of DAMA/LIBRA-phase2: **0.92 ton x yr**

Annual Cycles	Period	Mass (kg)	Exposure (kg · day)	( $\alpha-\beta^2$ )
1	Dec 2010 – Sept. 2011		Commissioning	
2	Nov. 2, 2011 – Sept. 11, 2012	242.5	62917	0.519
3	Oct. 8, 2012 – Sept. 2, 2013	242.5	60586	0.534
4	Sept. 8, 2013 – Sept. 1, 2014	242.5	73792	0.479
5	Sept. 1, 2014 – Sept. 9, 2015	242.5	71180	0.486
6	Sept. 10, 2015 – Aug. 24, 2016	242.5	67527	0.522
7	Sept 2016 –	242.5	running	

PRELIMINARY

Exposure expected for the first data release of DAMA/LIBRA-phase2, 6 a.c.:  $\approx 1.1$  ton x yr



# Towards future DAMA/LIBRA-phase3

**DAMA/LIBRA-phase3** (enhancing sensitivities for corollary aspects, other DM features, second order effects and other rare processes):

- R&D studies towards the possible DAMA/LIBRA-phase3 are continuing in particular as regards new protocols for possible modifications of the detectors; moreover, four new PMT prototypes from a dedicated R&D with HAMAMATSU are already at hand.
- Improving the light collection of the detectors (and accordingly the light yields and the energy thresholds). Improving the electronics.
- **Other possible option:** new ULB crystal scintillators (e.g.  $\text{ZnWO}_4$ ) placed in between the DAMA/LIBRA detectors to add also a high sensitivity directionality meas.

The presently-reached metallic PMTs features:

- Q.E. around 35-40% @ 420 nm (NaI(Tl) light)
- radiopurity at level of 5 mBq/PMT ( $^{40}\text{K}$ ), 3-4 mBq/PMT ( $^{232}\text{Th}$ ), 3-4 mBq/PMT ( $^{238}\text{U}$ ), 1 mBq/PMT ( $^{226}\text{Ra}$ ), 2 mBq/PMT ( $^{60}\text{Co}$ ).



4 prototypes at hand



## Other signatures?

- *Second order effects*
- *Diurnal effects*
- *Shadow effects*
- *Directionality*
- ...



# Diurnal effects

Velocity of the detector in the terrestrial laboratory:

Since:

$$|\vec{v}_s| = |\vec{v}_{LSR} + \vec{v}_\odot| \approx 232 \pm 50 \text{ km/s,}$$

$$|\vec{v}_{rev}(t)| \approx 30 \text{ km/s}$$

$$|\vec{v}_{rot}(t)| \approx 0.34 \text{ km/s} \quad \text{at LNGS}$$

$$v_{lab}(t) \simeq v_s + \hat{v}_s \cdot \vec{v}_{rev}(t) + \hat{v}_s \cdot \vec{v}_{rot}(t).$$

Expected signal counting rate in a given k-th energy bin:

$$S_k[v_{lab}(t)] \simeq S_k[v_s] + \left[ \frac{\partial S_k}{\partial v_{lab}} \right]_{v_s} [V_{Earth} B_m \cos \omega(t - t_0) + V_r B_d \cos \omega_{rot}(t - t_d)]$$

The ratio  $R_{dy}$  is a model independent constant:

$$R_{dy} = \frac{S_d}{S_m} = \frac{V_r B_d}{V_{Earth} B_m} \simeq 0.016 \quad \text{at LNGS latitude}$$

- Observed annual modulation amplitude in DAMA/LIBRA-phase1 in the (2–6) keV energy interval:  $(0.0097 \pm 0.0013) \text{ cpd/kg/keV}$
- Thus, the expected value of the diurnal modulation amplitude is  $\simeq 1.5 \times 10^{-4} \text{ cpd/kg/keV}$ .
- When fitting the *single-hit* residuals with a cosine function with amplitude  $A_d$  as free parameter, period fixed at 24 h and phase at 14 h: all the diurnal modulation amplitudes are compatible with zero.


$$A_d(2-6 \text{ keV}) < 1.2 \times 10^{-3} \text{ cpd/kg/keV (90\%CL)}$$




larger exposure DAMA/LIBRA-phase2 with lower energy threshold offers increased sensitivity to such an effect

A diurnal effect with the sidereal time is expected for DM because of Earth rotation

EPJC 74 (2014) 2827



Model-independent result on possible diurnal effect in DAMA/LIBRA-phase1




Present experimental sensitivity more modest than the expected diurnal modulation amplitude derived from the DAMA/LIBRA-phase1 observed effect.




# Earth shadowing effect with DAMA/LIBRA-phase1



EPJC75 (2015) 239



- **Earth Shadow Effect** could be expected for DM candidate particles inducing just nuclear recoils
- can be pointed out only for candidates with high cross-section with ordinary matter (low DM local density)
- would be induced by the variation during the day of the Earth thickness crossed by the DM particle in order to reach the experimental set-up



- DM particles crossing Earth lose their energy
- DM velocity distribution observed in the laboratory frame is modified as function of time (**GMST 8:00 black**; **GMST 20:00 red**)



Taking into account the DAMA/LIBRA DM annual modulation result, allowed regions in the  $\xi$  vs  $\sigma_n$  plane for each  $m_{DM}$ .

# Development of detectors with anisotropic response


DAMA - Seminal paper: N.Cim.C15(1992)475; revisited: EPJC28(2003)203; more recently more suitable materials: Eur. Phys. J. C 73 (2013) 2276; now: work in progress

Anisotropic detectors are of great interest for many applicative fields, e.g.:

⇒ they can offer a unique way to study directionality for Dark Matter candidates that induce just nuclear recoils

Taking into account:

- the correlation between the direction of the nuclear recoils and the Earth motion in the galactic rest frame;
- the peculiar features of anisotropic detectors;




The detector response is expected to vary as a function of the sidereal time

## Development of $\text{ZnWO}_4$ scintillators

- ✓ Both light output and pulse shape have anisotropic behavior and can provide two independent ways to study directionality
- ✓ Very high reachable radio-purity;
- ✓ Threshold at keV feasible;

O → light masses  
Zn, W → high masses



Presently running at ENEA-Casaccia  
with neutron generator to measure anisotropy  
in keV range



# $\text{ZnWO}_4$ – work in progress...

- ❖ Cryostat for low temperature measurement with scintillation detectors realized
- ❖ Test of the Cryostat in progress
- ❖ Lowering the energy threshold (new PMT with higher QE, SiPM, APD, SDD, ...)
- ❖ New purification techniques under study



- ❖ Measurements of anisotropy at low energy with MP320 Neutron Generator ( $E_n = 14 \text{ MeV}$ ) in progress at Casaccia ENEA lab
- ❖ Development of electronics

Exp @ ENEA-Casaccia lab



PSD capability of the EJ-309 Liquid Scintillator Detector Used



# Conclusions

- Positive evidence for the presence of DM particles in the galactic halo at  $9.3\sigma$  C.L. (14 annual cycles DAMA/NaI and DAMA/LIBRA-phase1: **1.33 ton  $\times$  yr**)
- Modulation parameters determined with higher precision
- New investigations on different peculiarities of the DM signal exploited (**Diurnal Modulation** and **Earth Shadow Effect**)
- New corollary analysis on **Mirror Dark Matter**
- Full sensitivity to many kinds of DM candidates and interactions types (both inducing recoils and/or e.m. radiation), **full sensitivity to low and high mass candidates**



- **DAMA/LIBRA – phase2** in **data taking** at lower software energy threshold (below 2 keV)
- Continuing investigations of rare processes other than DM
- **DAMA/LIBRA – phase3 R&D in progress**
- R&D for a possible DAMA/1ton set-up, proposed by DAMA since 1996, **continuing** as well as **some other R&Ds**

