

Solar neutrinos in dark matter detectors

Neutrinos and dark matter in nuclear physics (NDM 2018)



Daejeon, Korea

June 29-July 4, 2018

Louis E. Strigari



Neutrino backgrounds to dark matter detection



Neutrino-nucleus coherent scattering

Sensitive to BSM physics:

NSI: Scholberg 2005; Barranco et al. 2007


Sterile neutrinos: Dutta et al. 1508.07981, 1511.02834

Z' interactions: Lindner et al 2017; Abdullah et al. 2018

Nuclear physics: Li talk

$$\frac{d\sigma_{CNS}(E_\nu, T_R)}{dT_R} = \frac{G_f^2}{4\pi} Q_w^2 m_N \left(1 - \frac{m_N T_R}{2E_\nu^2}\right) F^2(T_R)$$

About a year ago “...a well known prediction of the Standard Model, but is yet to be detected....”






Brice et al, 1311.5958

COHERENT collaboration, Science, 2017



SNS flux (1.4 MW): $430 \times 10^5 \nu/\text{cm}^2/\text{s}$ @ 20 m;
~400 ns proton pulses @ 60 Hz \rightarrow $\sim 10^{-4}$ bg rejection



G. Rich talk

Coherent neutrino scattering at reactors

The CONNIE experiment

A. Aguilar-Arevalo¹, X. Bertou², C. Bonifazi³, M. Butner⁴,
G. Cancelo⁴, A. Castaneda Vazquez¹, B. Cervantes Vergara¹,
C.R. Chavez⁵, H. Da Motta⁶, J.C. D'Olive¹, J. Dos Anjos⁶,
J. Estrada⁴, G. Fernandez Moroni^{7,8}, R. Ford⁴, A. Foguel^{3,6},
K.P. Hernandez Torres¹, F. Izraelevitch⁴, A. Kavner⁹,
B. Kilminster¹⁰, K. Kuk⁴, H.P. Lima Jr.⁶, M. Makler⁶, J. Molina⁵,
G. Moreno-Granados¹, J.M. Moro¹¹, E.E. Paolini^{7,12}, M. Sofo Haro²,
J. Tiffenberg⁴, F. Trillaud¹, and S. Wagner^{6,13}

M. Lindner talk; CONUS
R. Strauss talk: Nu-cleus

Coherent Neutrino Scattering with Low Temperature Bolometers at Chooz Reactor Complex

J. Billard¹, R. Carr², J. Dawson³, E. Figueroa-Feliciano⁴, J. A.
Formaggio², J. Gascon¹, M. De Jesus¹, J. Johnston², T.
Lasserre^{5,6}, A. Leder², K. J. Palladino⁷, S. H. Trowbridge², M.
Vivier⁵, and L. Winslow²

Research program towards observation of neutrino-nucleus coherent scattering

H T Wong^{1,*}, H B Li¹, S K Lin¹, S T Lin¹, D He², J Li², X Li², Q
Yue², Z Y Zhou³ and S K Kim⁴

¹ Institute of Physics, Academia Sinica, Taipei 11529, Taiwan.


² Department of Engineering Physics, Tsing Hua University, Beijing 100084, China.


³ Department of Nuclear Physics, Institute of Atomic Energy, Beijing 102413, China.

⁴ Department of Physics, Seoul National University, Seoul 151-742, Korea.

Background Studies for the MINER Coherent Neutrino Scattering Reactor Experiment

G. Agnolet^a, W. Baker^a, D. Barker^b, R. Beck^a, T.J. Carroll^c, J. Cesar^c, P. Cushman^b, J.B. Dent^d,
S. De Rijck^c, B. Dutta^a, W. Flanagan^c, M. Fritts^b, Y. Gao^{a,e}, H.R. Harris^a, C.C. Hays^a, V. Iyer^f,
A. Jastram^a, F. Kadribasic^a, A. Kennedy^b, A. Kubik^a, I. Ogawa^g, K. Lang^c, R. Mahapatra^a, V. Mandic^b,
R.D. Martin^h, N. Mast^b, S. McDeavittⁱ, N. Mirabolfathi^a, B. Mohanty^f, K. Nakajima^g, J. Newhouseⁱ,
J.L. Newstead^j, D. Phan^c, M. Proga^c, A. Roberts^k, G. Rogachev^l, R. Salazar^c, J. Sander^k, K. Senapati^f,
M. Shimada^g, L. Strigari^a, Y. Tamagawa^g, W. Teizer^a, J.I.C. Vermaakⁱ, A.N. Villano^b, J. Walker^m,
B. Webb^a, Z. Wetzel^a, S.A. Yadavalli^c







Reactors



Accelerators





Reactors



Accelerators





Distinguishing DM from neutrinos:

- Annual modulation/
directionality: Grothaus,
Fairbairn, Monroe 2014; O'Hare
et al. 2015, Davis 2015
- SD DM: Ruppin, Billard,
Figueroa-Feliciano, Strigari,
2014; Gelmini et al. 2018
- Non-rel EFTs: Dent, Dutta,
Strigari, Newstead 2016/2017
- NSI: Dutta, Liao, LS, Walker
2017; Aristizabal Sierra, Rojas,
Tytgat 2018; Gonzalez-Garcia et
al. 2018



Aristizabal Sierra talk

Neutrino floor for light DM



- At low mass, neutrino floor from solar neutrinos
- Particularly important for detectors that lack electron/nuclear discrimination

Solar neutrinos: Status



Solar Neutrinos: Status and Prospects

W.C. Haxton,¹ R.G. Hamish Robertson,²
and Aldo M. Serenelli³

The program of solar neutrino studies envisioned by Davis and Bahcall has been only partially completed. Borexino has extended precision measurements

to low-energy solar neutrinos, determining the flux of ^7Be neutrinos to 5%, and thereby confirming the expected increase in the ν_e survival probability for neutrino energies in the vacuum-dominated region. First results on the pep neutrino

High-Z Low-Z

| ν flux | E_ν^{\max} (MeV) | GS98-SFII | AGSS09-SFII | Solar | units |
|---|----------------------|---------------------|---------------------|-----------------------------|-------------------------------|
| $p+p \rightarrow {}^2\text{H}+e^++\nu$ | 0.42 | $5.98(1 \pm 0.006)$ | $6.03(1 \pm 0.006)$ | $6.05(1^{+0.003}_{-0.011})$ | $10^{10}/\text{cm}^2\text{s}$ |
| $p+e^-+p \rightarrow {}^2\text{H}+\nu$ | 1.44 | $1.44(1 \pm 0.012)$ | $1.47(1 \pm 0.012)$ | $1.46(1^{+0.010}_{-0.014})$ | $10^8/\text{cm}^2\text{s}$ |
| ${}^7\text{Be}+e^- \rightarrow {}^7\text{Li}+\nu$ | 0.86 (90%) | $5.00(1 \pm 0.07)$ | $4.56(1 \pm 0.07)$ | $4.82(1^{+0.05}_{-0.04})$ | $10^9/\text{cm}^2\text{s}$ |
| | 0.38 (10%) | | | | |
| ${}^8\text{B} \rightarrow {}^8\text{Be}+e^++\nu$ | ~ 15 | $5.58(1 \pm 0.14)$ | $4.59(1 \pm 0.14)$ | $5.00(1 \pm 0.03)$ | $10^6/\text{cm}^2\text{s}$ |
| ${}^3\text{He}+p \rightarrow {}^4\text{He}+e^++\nu$ | 18.77 | $8.04(1 \pm 0.30)$ | $8.31(1 \pm 0.30)$ | — | $10^3/\text{cm}^2\text{s}$ |
| ${}^{13}\text{N} \rightarrow {}^{13}\text{C}+e^++\nu$ | 1.20 | $2.96(1 \pm 0.14)$ | $2.17(1 \pm 0.14)$ | ≤ 6.7 | $10^8/\text{cm}^2\text{s}$ |
| ${}^{15}\text{O} \rightarrow {}^{15}\text{N}+e^++\nu$ | 1.73 | $2.23(1 \pm 0.15)$ | $1.56(1 \pm 0.15)$ | ≤ 3.2 | $10^8/\text{cm}^2\text{s}$ |
| ${}^{17}\text{F} \rightarrow {}^{17}\text{O}+e^++\nu$ | 1.74 | $5.52(1 \pm 0.17)$ | $3.40(1 \pm 0.16)$ | $\leq 59.$ | $10^6/\text{cm}^2\text{s}$ |
| χ^2/P^{agr} | | 3.5/90% | 3.4/90% | | |

Haxton et al. 2013

- 3D rotational hydrodynamical simulations suggest lower metallicity in Solar core (Asplund et al. 2009)
- Low metallicity in conflict with heliosiesmology data
- SNO Neutral Current measurement right in between predictions of low and high metallicity SSMs

Serenelli talk


High-Z Low-Z

| ν flux | E_ν^{\max} (MeV) | GS98-SFII | AGSS09-SFII | Solar | units |
|---|----------------------|---------------------|---------------------|-----------------------------|-------------------------------|
| $p+p \rightarrow {}^2\text{H}+e^++\nu$ | 0.42 | $5.98(1 \pm 0.006)$ | $6.03(1 \pm 0.006)$ | $6.05(1^{+0.003}_{-0.011})$ | $10^{10}/\text{cm}^2\text{s}$ |
| $p+e^-+p \rightarrow {}^2\text{H}+\nu$ | 1.44 | $1.44(1 \pm 0.012)$ | $1.47(1 \pm 0.012)$ | $1.46(1^{+0.010}_{-0.014})$ | $10^8/\text{cm}^2\text{s}$ |
| ${}^7\text{Be}+e^- \rightarrow {}^7\text{Li}+\nu$ | 0.86 (90%) | $5.00(1 \pm 0.07)$ | $4.56(1 \pm 0.07)$ | $4.82(1^{+0.05}_{-0.04})$ | $10^9/\text{cm}^2\text{s}$ |
| | 0.38 (10%) | | | | |
| ${}^8\text{B} \rightarrow {}^8\text{Be}+e^++\nu$ | ~ 15 | $5.58(1 \pm 0.14)$ | $4.59(1 \pm 0.14)$ | $5.00(1 \pm 0.03)$ | $10^6/\text{cm}^2\text{s}$ |
| ${}^3\text{He}+p \rightarrow {}^4\text{He}+e^++\nu$ | 18.77 | $8.04(1 \pm 0.30)$ | $8.31(1 \pm 0.30)$ | — | $10^3/\text{cm}^2\text{s}$ |
| ${}^{13}\text{N} \rightarrow {}^{13}\text{C}+e^++\nu$ | 1.20 | $2.96(1 \pm 0.14)$ | $2.17(1 \pm 0.14)$ | ≤ 6.7 | $10^8/\text{cm}^2\text{s}$ |
| ${}^{15}\text{O} \rightarrow {}^{15}\text{N}+e^++\nu$ | 1.73 | $2.23(1 \pm 0.15)$ | $1.56(1 \pm 0.15)$ | ≤ 3.2 | $10^8/\text{cm}^2\text{s}$ |
| ${}^{17}\text{F} \rightarrow {}^{17}\text{O}+e^++\nu$ | 1.74 | $5.52(1 \pm 0.17)$ | $3.40(1 \pm 0.16)$ | $\leq 59.$ | $10^6/\text{cm}^2\text{s}$ |
| χ^2/P^{agr} | | 3.5/90% | 3.4/90% | | |

Haxton et al. 2013

- 3D rotational hydrodynamical simulations suggest lower metallicity in Solar core (Asplund et al. 2009)
- Low metallicity in conflict with heliosiesmology data
- SNO Neutral Current measurement right in between predictions of low and high metallicity SSMs



Serenelli talk



Borexino collaboration


- Borexino, SNO, SK indicate the low energy ES data lower than MSW predicts
- Upturn in MSW survival probability not been measured
- May indicate new physics (e.g. Holanda & Smirnov 2011)

Nuclear and electron recoil spectra



Non-Standard Neutrino Interactions


- NSI describe new physics at high energy in form of heavy scalars, gauge bosons
- Best sensitivity to flavor-conserving Neutral Current NSI models
- NSI identified in CNS detection



$$\mathcal{L}_{int} = 2\sqrt{2}G_F \bar{\nu}_{\alpha L} \gamma^\mu \nu_{\beta L} \left(\epsilon_{\alpha\beta}^{fL} \bar{f}_L \gamma_\mu f_L + \epsilon_{\alpha\beta}^{fR} \bar{f}_R \gamma_\mu f_R \right)$$


$$\frac{d\sigma}{dE_r} = \frac{2}{\pi} G_F^2 m_f \left[\left| \epsilon_{\alpha\beta}^{fL} \right|^2 + \left| \epsilon_{\alpha\beta}^{fR} \right|^2 \left(1 - \frac{E_r}{E_\nu} \right)^2 - \frac{1}{2} \left(\epsilon_{\alpha\beta}^{fL*} \epsilon_{\alpha\beta}^{fR} + \epsilon_{\alpha\beta}^{fL} \epsilon_{\alpha\beta}^{fR*} \right) \frac{m_f E_r}{E_\nu^2} \right]$$

Non-standard interactions + MSW + DM detectors




Friedland, Lunardini, Pena-Garay PLB 2004

Non-standard interactions + MSW + DM detectors





Friedland, Lunardini, Pena-Garay PLB 2004



B. Dutta, **Shu Liao**, L. Strigari, J. Walker, PLB 2017

- NSI may increase or decrease event rate in Xenon
- 1t sensitive to models still consistent with nu oscillations

Non-standard interactions + DM detectors





Carter Hall LZ talk

B. Dutta, **Shu Liao**, L. Strigari, J. Walker, PLB 2017

- NSI may increase or decrease event rate in Xenon
- 1t sensitive to models still consistent with nu oscillations


Low energy solar neutrino spectroscopy



- Multicomponent spectral analysis of low energy solar neutrinos
- 2.7% precision on ${}^7\text{Be}$
- Strongest upper bound on CNO neutrinos

CNO Solar neutrinos and neutrino luminosity

Future low energy neutrino electron elastic scattering experiments for CNO



CNO Solar neutrinos and neutrino luminosity

- Since nuclear fusion is dominant energy source, linear combination of neutrino fluxes equals the photon luminosity
- Deviation between *neutrino luminosity* and photon luminosity could hint at alternative sources of energy generation

$$\frac{L_{\text{pp-chain}}}{L_{\odot}} = 0.991^{+0.005}_{-0.004} \left[{}^{+0.008}_{-0.013} \right] \iff \frac{L_{\text{CNO}}}{L_{\odot}} = 0.009^{+0.004}_{-0.005} \left[{}^{+0.013}_{-0.008} \right]$$



$$\frac{L_{\odot}(\text{neutrino-inferred})}{L_{\odot}} = 1.04^{+0.07}_{-0.08} \left[{}^{+0.20}_{-0.18} \right]$$

Bergstrom, Gonzalez-Garcia et al.
JHEP 2016

- Since nuclear fusion is dominant energy source, linear combination of neutrino fluxes equals the photon luminosity


Direct pp measurement with Xe at few percent level can improve this constraint

CNO Solar neutrinos and neutrino luminosity



G3 Xe detector may be used for CNO (Newstead, LS, Lang)
Requires reduction of detector backgrounds




CNO Solar neutrinos and neutrino luminosity



G3 Xe detector may be used for CNO (Newstead, LS, Lang)
Requires reduction of detector backgrounds

eV-scale sterile neutrinos

- Combined with 'reactor anomaly', gallium results may hint at new physics, i.e. \sim eV sterile neutrino (Giunti & Laveder 2010; Mention 2011)



G3 detector can provide a test of the reactor/
gallium anomaly (Billard, LS, Figueroa-Feliciano,
PRD 2014, 1409.0050)

Recap: Neutrinos in dark matter experiments

Astrophysics

- First measurement of the 8B neutral current energy spectrum
- First direct measurement of the survival probability for low energy solar neutrinos
- Direct measurement of the CNO flux
- PP flux measurement to ~ few percent will provide most stringent measurement of the “neutrino luminosity” of the Sun

Recap: Neutrinos in dark matter experiments




Astrophysics

- First measurement of the 8B neutral current energy spectrum
- First direct measurement of the survival probability for low energy solar neutrinos
- Direct measurement of the CNO flux
- PP flux measurement to \sim few percent will provide most stringent measurement of the “neutrino luminosity” of the Sun




Particle physics

- NSI affects both neutrino-coherent scattering and neutrino-electron elastic scattering channels
- Independent probe of eV-scale sterile neutrinos

Reactor, accelerator, solar complementarity







Solar neutrinos add
sensitivity to NSI from
neutrino propagation



Dent, Dutta, Liao,
Newstead, LS,
Walker PRD 2018

New directions in dark matter and neutrino physics

Astrophysical sources



Reactors



Accelerators

