

# Modeling Evolution of Dark Matter Substructure and Annihilation Boost

Reference: Phys.Rev.D97., 123002

**Nagisa Hiroshima (Univ. of Tokyo, KEK)**

Collaboration with

S.Ando (Univ. of Amsterdam), T.Ishiyama (Chiba Univ.)

# Contents

- **Motivations**
- **Modeling**
- **Applications**
- **Summary**

# Motivations


# WIMP Dark Matter

- Naturally explains the relic abundance with weak scale cross section


$$\langle\sigma v\rangle \sim 3 \times 10^{-26} \text{cm}^3 \text{s}^{-1}$$

- WIMP DM mass should be  $m_{\text{DM}} \sim \mathcal{O}(1)\text{GeV} - \mathcal{O}(1)\text{TeV}$

- predict formation of small scale structures down to  $10^{-12} - 10^{-3} M_{\odot}$




# DM search with $\gamma$ -rays



$$\phi_\gamma = \frac{1}{2} \frac{1}{4\pi} \frac{\langle \sigma v \rangle}{m_{\text{DM}}^2} \int \frac{dN_\gamma}{dE_\gamma} dE_\gamma \cdot$$

$$\int_{l.o.s} \rho_{\text{DM}}^2 ds$$

# subhalo boost



$$\phi_\gamma = \frac{1}{2} \frac{1}{4\pi} \frac{\langle \sigma v \rangle}{m_{\text{DM}}^2} \int \frac{dN_\gamma}{dE_\gamma} dE_\gamma \cdot (1+B) \int_{l.o.s} \rho_{\text{DM}}^2 ds$$


**subhalos on l.o.s boost the annihilation signals**

# Estimates of subhalo boost

1312.1729

- ♦  $M_{\text{halo}} \sim 10^{-6} - 10^{16} M_{\odot}$
- ♦  $z \sim 0 - 10$

**Previous works adopted  
extrapolation of results from  
numerical simulations**



**Needs for physical, wide-coverage modelings**

# Modeling



# Evolution of subhalos


1. Formation

2. Accretion

3. Tidal Stripping


Assumptions:

- NFW profile with truncation for host & subhalo
- mass-loss occurs in the first orbit of each subhalo



$$\dot{m} = \frac{m - m(r_t)}{T_r}$$

# Tidal Stripping



a single power-law in 20 orders of magnitude


# calculations of boost factors

- ✦ **host evolutions**
- ✦ **mass accretion history**
- ✦ **NFW parameters after tidal stripping**

**Boost factor**  $B = \frac{\sum_i w_i \rho_{s,i}^2 r_{s,i}^3}{\rho_{s,\text{host}}^2 r_{s,\text{host}}^3}$


# Applications

# Mass fraction of subhalo



Good agreements with N-body result


# Boost Factor



**Annihilation signal is boosted by factors**

# Update on IGRB limit

**DM + DM  $\rightarrow$  b  $\bar{b}$**



**Improving IGRB constraint**

# Summary




# Summary

- **We modeled tidal stripping of subhalos in an analytical way covering more than 20 orders of mass ranges**
- **Subhalos in our model show good agreements with those in N-body simulations**
- **As applications, we calculate boost factor of DM annihilation signals**
- **DM annihilation signals can be boosted up to a factor of 10 in cluster scales**



# Boost Factor



Annihilation signal is boosted by factors

# Tidal stripping

- **potential of the host**

$$\Phi(R) = -\frac{GM_{\text{host}}}{R_{\text{vir}}} \frac{\ln \left[ 1 + c_{\text{vir}}^{\text{host}} R/R_{\text{vir}} \right]}{f(c_{\text{vir}})^{\text{host}} R/R_{\text{vir}}}$$

- **orbital period of subhalo**


$$T_r = 2 \int_{R_p}^{R_a} \frac{dR}{\sqrt{2 \left[ E - \Phi(R) \right] - L^2/R^2}}$$

- **tidal mass of subhalo**


$$(r_t/R_p)^3 = \frac{m(r_t)}{M_{<R_p}} \left( 2 + \left( L^2/R_p GM_{<R_p} \right) - \left( d \ln M(R)/d \ln R \right) |_{R_p} \right)^{-1}$$

- **mass-loss rate**

$$\dot{m} = \left[ m - m(r_t) \right] / T_r$$



# Estimates of subhalo boost



**Needs for analytical modelings**