

Galileon and generalized Galileon with projective invariance in metric-affine formalism

Katsuki Aoki, Waseda University

with K. Shimada

Based on Phys.Rev. D98 (2018) 044038 (arXiv: 1806.02589).

Introduction

GR is 1) a theory of massless spin-2 field
2) a theory of curved geometry

Extensions of GR (focusing on 1?)

Massless → massive: massive gravity, bigravity...

Spin-2 → other spin: scalar-tensor, vector-tensor...

We have to take care 2 when extending GR.

Physics should require how to measure the distance and the derivative.

→ **two independent objects, metric $g_{\mu\nu}$ and connection $\Gamma_{\alpha\beta}^{\mu}$.**

Riemannian geometry: metric is only independent object

$$\Gamma_{\alpha\beta}^{\mu} = \left\{ \begin{matrix} \mu \\ \alpha\beta \end{matrix} \right\} := \frac{1}{2} g^{\mu\nu} (\partial_{\alpha} g_{\beta\nu} + \partial_{\beta} g_{\alpha\nu} - \partial_{\nu} g_{\alpha\beta})$$

Just a special case!

Metric and Metric-affine formalisms

- **Metric formalism:** Gravity is a theory of metric (= spin-2 field)
→ Gravitational theory determines only metric (Riemannian geometry)
- **Metric-affine (Palatini) formalism:** Gravity is a theory of geometry
→ **Gravitational theory determines not only metric but also connection.**
No assumption on the connection.

GR in metric formalism is obtained from GR in metric-affine formalism.
(Giachetta and Mangiarotti 1997, Dadhich and Pons, 2012 for example)

$$S_{\text{gravity}}(g, \Gamma) = \int d^4x \sqrt{-g} \frac{M_{\text{pl}}^2}{2} \overset{\Gamma}{R}(g, \Gamma) + \text{higher curvatures}$$

EH term

$$\overset{\Gamma}{R}{}^{\mu}{}_{\nu\alpha\beta}(\Gamma) := \partial_{\alpha}\Gamma_{\beta\nu}^{\mu} - \partial_{\beta}\Gamma_{\alpha\nu}^{\mu} + \Gamma_{\alpha\sigma}^{\mu}\Gamma_{\beta\nu}^{\sigma} - \Gamma_{\beta\sigma}^{\mu}\Gamma_{\alpha\nu}^{\sigma}, \quad \overset{\Gamma}{R} := g^{\mu\nu}\overset{\Gamma}{R}_{\mu\nu}, \quad \overset{\Gamma}{R}_{\mu\nu} := \overset{\Gamma}{R}{}^{\alpha}{}_{\mu\alpha\beta}$$

Metric-affine formalism

For convenience, we introduce the distortion tensor κ

$$\kappa^\mu{}_{\alpha\beta} := \Gamma^\mu{}_{\alpha\beta} - \left\{ \begin{matrix} \mu \\ \alpha\beta \end{matrix} \right\}$$

The metric-affine formalism = The metric formalism with κ .

$${}^\Gamma R^\mu{}_{\alpha\beta\gamma}(\Gamma) = R^\mu{}_{\alpha\beta\gamma}(g) + 2\nabla_{[\alpha}\kappa^\mu{}_{\beta]\nu} + \kappa^\mu{}_{[\alpha|\sigma}\kappa^\sigma{}_{\beta]\nu}$$

$$\mathcal{L}_{\text{EH}} \sim M_{\text{pl}}^2 R(g) + M_{\text{pl}}^2 \kappa^2 : \text{mass term of distortion}$$

$$\text{higher curvature} \supset (\nabla\kappa)^2 : \text{kinetic term of distortion}$$

When higher curvatures can be ignored, κ can be integrated out.

$$\Rightarrow \kappa^\mu{}_{\alpha\beta} = 0 : \text{Riemannian geometry} \quad \Gamma^\mu{}_{\alpha\beta} = \left\{ \begin{matrix} \mu \\ \alpha\beta \end{matrix} \right\} := \frac{1}{2}g^{\mu\nu}(\partial_\alpha g_{\beta\nu} + \partial_\beta g_{\alpha\nu} - \partial_\nu g_{\alpha\beta})$$

We **don't need to assume anything** on the connection to get GR.

Metric or Metric-affine?

- Einstein gravity: Riemannian = beyond Riemannian
- Beyond Einstein: Riemannian \neq beyond Riemannian

For example, higher curvature corrections

$$\text{e.g., } f(R), f(R_{\mu\nu})$$

We consider scalar-tensor theories with higher derivatives.

$$\nabla^{\Gamma} \nabla^{\Gamma} \phi \supset \kappa \partial \phi$$

- ✓ Uniqueness of Galileon
- ✓ Simple reformulation of DHOST (no fine-tuning)
- ✓ Scalar-fermion non-minimal coupling

A point: metric-affine formalism can have an additional symmetry.

Projective invariance

$$\mathcal{L}_{\text{EH}}(g, \Gamma) = \frac{M_{\text{pl}}^2}{2} R = \frac{M_{\text{pl}}^2}{2} (R(g) + \kappa^\alpha_{\alpha\beta} \kappa^{\beta\gamma}_{\gamma} - \kappa^{\alpha\beta}_{\gamma} \kappa^{\gamma}_{\alpha\beta}) + \text{total divergence}$$

The EH action has an additional symmetry, "**projective invariance**",

$$\Gamma^\mu_{\alpha\beta} \rightarrow \Gamma^\mu_{\alpha\beta} + \delta^\mu_{\beta} U_{\alpha}(x) \quad \text{or} \quad \kappa^\mu_{\alpha\beta} \rightarrow \kappa^\mu_{\alpha\beta} + \delta^\mu_{\beta} U_{\alpha}(x)$$

which preserves the geodesic equation

$$\frac{d^2 x^\mu}{d\lambda^2} + \Gamma^\mu_{\alpha\beta} \frac{dx^\alpha}{d\lambda} \frac{dx^\beta}{d\lambda} = 0$$

and the angle for the parallel transport (a kind of conformal symmetry)

Also, all standard matter Lagrangians are projective invariant.

Let's assume Galileon is projective invariant → Uniqueness of Galileon

Galileon scalar field

- Flat spacetime (Euclidean geometry): we have only one Galileon

$$\mathcal{L}_n^{\text{gal}} = \epsilon \epsilon (\partial\phi)^2 (\partial\partial\phi)^{n-2} = (\partial\phi)^2 \epsilon \epsilon (\partial\partial\phi)^{n-2} + \text{total divergence}$$

$$\begin{aligned} \text{e.g., } \mathcal{L}_4^{\text{gal}} &= \epsilon^{\alpha\beta\gamma\delta} \epsilon^{\alpha'\beta'\gamma'} \delta_{\alpha\beta\gamma\delta} \partial_{\alpha'} \phi \partial_{\alpha'} \phi \partial_{\beta'} \phi \partial_{\beta'} \phi \partial_{\gamma'} \phi \partial_{\gamma'} \phi \\ &= \partial_{\mu} \phi \partial^{\mu} \phi \epsilon^{\alpha\beta\gamma\delta} \epsilon^{\alpha'\beta'} \gamma_{\delta} \partial_{\alpha'} \partial_{\alpha} \phi \partial_{\beta'} \partial_{\beta} \phi + \text{total divergence} \end{aligned}$$

Two are same via integration by parts.

- Curved spacetime (Riemannian geometry): we have **two** Galileons

$$\epsilon \epsilon (\nabla\phi)^2 (\nabla\nabla\phi)^{n-2} \neq (\nabla\phi)^2 \epsilon \epsilon (\nabla\nabla\phi)^{n-2} + \text{total divergence}$$

GLPV (covariantized)

Horndeski (covariant)

Two are **not** same.

Galileon with projective invariance

- Curved spacetime (**metric-affine** geometry):

Due to the projective invariance Galileon must be

$$\mathcal{L}_n^{\text{gal}\Gamma}(\overset{\Gamma}{\nabla}_\mu\phi, \overset{\Gamma}{\nabla}_\mu\overset{\Gamma}{\nabla}_\nu\phi; g_{\mu\nu}) = \epsilon\epsilon(\overset{\Gamma}{\nabla}\phi)^2(\overset{\Gamma}{\nabla}\overset{\Gamma}{\nabla}\phi)^{n-2} \quad (\text{GLPV type})$$

$$\Rightarrow \mathcal{L}(g, \Gamma, \phi) = \frac{M_{\text{pl}}^2}{2} g^{\mu\nu} R_{\mu\nu}^\Gamma + \sum_{n \geq 2}^5 \frac{c_n}{\Lambda^{3(n-2)}} \mathcal{L}_n^{\text{gal}\Gamma}$$

$$\text{with } \mathcal{L}_n^{\text{gal}\Gamma} = \epsilon\epsilon(\overset{\Gamma}{\nabla}\phi)^2(\overset{\Gamma}{\nabla}\overset{\Gamma}{\nabla}\phi)^{n-2} \supset \kappa^{n-2}(\partial\phi)^n$$

Up to quartic order ($n \leq 4$) $\rightarrow \kappa$ can be explicitly solved.

(The equation of κ becomes nonlinear when including quintic order)

After integrating out κ , we obtain...

Galileon with projective invariance

Effective action in Riemannian geometry

$$\mathcal{L}(g, \Gamma(g, \phi), \phi) = \frac{M_{\text{pl}}^2}{2} R(g) + \frac{3(c_3^2 - 4c_2c_4)X^3/\Lambda_2^8}{1 + 2c_4X^2/\Lambda_2^8} + \frac{1}{1 + 2c_4X^2/\Lambda_2^8} \left(c_2\mathcal{L}_2^{\text{galg}} + \frac{c_3}{\Lambda_3^3}\mathcal{L}_3^{\text{galg}} + \frac{c_4}{\Lambda_3^6}\mathcal{L}_4^{\text{galg}} \right)$$

$$X = (\partial\phi)^2, \quad \mathcal{L}_n^{\text{galg}} = \epsilon\epsilon(\nabla\phi)^2(\nabla\nabla\phi)^{n-2}$$

$$\Lambda_2^4 = \Lambda_3^3 M_{\text{pl}}$$

- ✓ does not coincide with either covariantized or covariant Galileon.
- ✓ can yield the non-minimal coupling to the fermion current

$$\mathcal{L}_{\text{int}} = \frac{i}{M_{\text{pl}}^2(1 + 2c_4X^2/\Lambda_2^8)} \left[\frac{3c_3}{2\Lambda_3^3}X + \frac{c_4}{\Lambda_3^6} \left(X\phi_\beta^\beta - \phi^{\beta\gamma}\phi_\beta\phi_\gamma \right) \right] j_\alpha\phi^\alpha + \dots$$

$$\mathcal{L}_D = \frac{i}{2}\bar{\psi}\gamma^\mu D_\mu\psi \supset -\frac{1}{4}\epsilon^{\alpha\beta\gamma\delta}\kappa_{\alpha\beta\gamma}j_\delta^5 + \frac{i}{2}\kappa^{[\alpha\beta]}{}_\beta j_\alpha, \quad j_\alpha = \bar{\psi}\gamma_\alpha\psi, \quad \phi_\mu = \nabla_\mu\phi,$$

$$j_\alpha^5 = \bar{\psi}\gamma_\alpha\gamma^5\psi, \quad \phi_{\mu\nu} = \nabla_\mu\nabla_\nu\phi$$

Generalized Galileon is DHOST

In metric formalism, generalizations of Galileon = Horndeski, GLPV

A straightforward generalization (up to quadratic in connection)

$$\begin{aligned}\mathcal{L}(g, \Gamma, \phi) &= f_1(\phi, X) \overset{\Gamma}{R} + f_2(\phi, X) \overset{\Gamma}{G}^{\mu\nu} \overset{\Gamma}{\nabla}_\mu \phi \overset{\Gamma}{\nabla}_\nu \phi \\ &+ F_2(\phi, X) + F_3(\phi, X) \mathcal{L}_3^{\text{gal}\Gamma} + F_4(\phi, X) \mathcal{L}_4^{\text{gal}\Gamma}\end{aligned}$$

= Non-minimal couplings + generalized Galileon

where Ricci scalar $\overset{\Gamma}{R} := g^{\mu\nu} \overset{\Gamma}{R}_{\mu\nu}$, $\overset{\Gamma}{R}_{\mu\nu} := \overset{\Gamma}{R}{}^{\alpha}{}_{\mu\alpha\nu}$ $X := g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi$

Einstein tensor $\overset{\Gamma}{G}^{\alpha\beta} := \frac{1}{4} \epsilon^{\gamma\alpha\mu\nu} \epsilon_{\gamma}{}^{\beta\mu'\nu'} \overset{\Gamma}{R}_{\mu\nu\mu'\nu'}$

Need fine-tuning of functions to be ghost-free?

→ Don't need! This action yields **class N-1/Ia of DHOST = ghost-free**

Metric formalism of DHOST

$$\mathcal{L}(g, \phi) = fR(g) + P + Q_1 g^{\mu\nu} \phi_{\mu\nu} + Q_2 \phi^\mu \phi_{\mu\nu} \phi^\nu + C^{\mu\nu, \rho\sigma} \phi_{\mu\nu} \phi_{\rho\sigma}$$

$$C^{\mu\nu, \rho\sigma} = \alpha_1 g^{\rho(\mu} g^{\nu)\sigma} + \alpha_2 g^{\mu\nu} g^{\rho\sigma} + \frac{1}{2} \alpha_3 (\phi^\mu \phi^\nu g^{\rho\sigma} + \phi^\rho \phi^\sigma g^{\mu\nu}) \\ + \frac{1}{2} \alpha_4 (\phi^\rho \phi^{(\mu} g^{\nu)\sigma} + \phi^\sigma \phi^{(\mu} g^{\nu)\rho}) + \alpha_5 \phi^\mu \phi^\nu \phi^\rho \phi^\sigma$$

DHOST is defined by $D_0 = D_1 = D_2 = 0$

$$f = f(\phi, X), \alpha_i = \alpha_i(\phi, X)$$

$$D_0 := -4(\alpha_1 + \alpha_2)[Xf(2\alpha_1 + X\alpha_6 + 4f_X) - 2f^2 - 8X^2 f_X^2],$$

$$D_1 := 4[X^2 \alpha_1(\alpha_1 + 3\alpha_2) - 2f^2 - 4Xf\alpha_2]\alpha_4 + 4X^2 f(\alpha_1 + \alpha_2)\alpha_5 + 8X\alpha_1^3 \\ - 4(f + 4Xf_X - 6X\alpha_2)\alpha_1^2 - 16(f + 5Xf_X)\alpha_1\alpha_2 \\ + 4X(3f - 4Xf_X)\alpha_1\alpha_3 - X^2 f\alpha_3^2 \\ + 32f_X(f + 2Xf_X)\alpha_2 - 16ff_X\alpha_1 - 8f(f - Xf_X)\alpha_3 + 48ff_X^2,$$

$$D_2 := 4[2f^2 + 4Xf\alpha_2 - X^2 \alpha_1(\alpha_1 + 3\alpha_2)]\alpha_5 + 4\alpha_1^3$$

$$+ 4(2\alpha_2 - X\alpha_3 - 4f_X)\alpha_1^2 + 3X^2 \alpha_1 \alpha_3^2 - 4Xf\alpha_3^2$$

$$+ 8(f + Xf_X)\alpha_1\alpha_3 - 32f_X\alpha_1\alpha_2 + 16f_X^2\alpha_1 + 32f_X^2\alpha_2 - 16ff_X\alpha_3$$

$$f_X = \partial f / dX$$

Metric-affine formalism of DHOST

$$\begin{aligned}\mathcal{L}(g, \Gamma, \phi) &= f_1(\phi, X) \overset{\Gamma}{R} + f_2(\phi, X) \overset{\Gamma}{G}{}^{\mu\nu} \overset{\Gamma}{\nabla}{}_{\mu} \phi \overset{\Gamma}{\nabla}{}_{\nu} \phi \\ &+ F_2(\phi, X) + F_3(\phi, X) \mathcal{L}_3^{\text{gal}\Gamma} + F_4(\phi, X) \mathcal{L}_4^{\text{gal}\Gamma}\end{aligned}$$

= Non-minimal couplings + generalized Galileon

in metric-affine formalism

Metric-affine formalism of DHOST

Integrating out κ ,

$$\mathcal{L}(g, \Gamma(g, \phi), \phi) = fR(g) + P + Q_1 g^{\mu\nu} \phi_{\mu\nu} + Q_2 \phi^\mu \phi_{\mu\nu} \phi^\nu + C^{\mu\nu, \rho\sigma} \phi_{\mu\nu} \phi_{\rho\sigma}$$

$$C^{\mu\nu, \rho\sigma} = \alpha_1 g^{\rho(\mu} g^{\nu)\sigma} + \alpha_2 g^{\mu\nu} g^{\rho\sigma} + \frac{1}{2} \alpha_3 (\phi^\mu \phi^\nu g^{\rho\sigma} + \phi^\rho \phi^\sigma g^{\mu\nu})$$

$$+ \frac{1}{2} \alpha_4 (\phi^\rho \phi^{(\mu} g^{\nu)\sigma} + \phi^\sigma \phi^{(\mu} g^{\nu)\rho}) + \alpha_5 \phi^\mu \phi^\nu \phi^\rho \phi^\sigma$$

where $f = f_1 - \frac{1}{2} f_2 X$, $P = F_2 + \frac{3X(f_1 \phi - F_3 X)^2}{2f_1 - f_2 X + 2F_4 X^2}$,

$$\phi_\mu = \nabla_\mu \phi, \phi_{\mu\nu} = \nabla_\mu \nabla_\nu \phi$$

$$Q_1 = -2f_\phi + \frac{4f_1(f_1 \phi - F_3 X)}{2f_1 - f_2 X + 2F_4 X^2}, \quad Q_2 = \frac{2f_\phi}{X} - \frac{4(f_1 - 3f_{1X})(f_1 \phi - F_3 X)}{X(2f_1 - f_2 X + 2F_4 X^2)},$$

$$\alpha_1 = -\alpha_2 = -\frac{f_2}{2} - \frac{f_1(f_2 - 2F_4 X)}{2f_1 - f_2 X + 2F_4 X^2}, \quad \alpha_3 = 2f_{2X} + \frac{4f_1 F_4 + (4f_{1X} - f_2)(f_2 - 2F_4 X)}{2f_1 - f_2 X + 2F_4 X^2},$$

$$\alpha_4 = -2f_{2X} + 2f_1^{-1} f_{1X}(3f_{1X} - f_2) + f_1^{-2} f_{1X} X(f_{1X} f_2 - 4f_1 f_{2X}) + \frac{f_2^2 - 4f_1 F_4 - 2f_2 F_4 X}{2f_1 - f_2 X + 2F_4 X^2},$$

$$\alpha_5 = -f_1^{-2} f_{1X}(f_{1X} f_2 - 4f_1 f_{2X}) + \frac{2f_{1X} \{4f_1 F_4 + (3f_{1X} - f_2)(f_2 - 2F_4 X)\}}{f_1(2f_1 - f_2 X + 2F_4 X^2)},$$

$$X := g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi$$

satisfy the degeneracy conditions $D_0 = D_1 = D_2 = 0$ (in class N-1/la)

Same number of arbitrary functions!

Specific model

We now know meaning of each function of DHOST.

$$\begin{aligned}\mathcal{L}(g, \Gamma, \phi) &= f_1(\phi, X) \overset{\Gamma}{R} + f_2(\phi, X) \overset{\Gamma}{G}{}^{\mu\nu} \overset{\Gamma}{\nabla}{}_{\mu} \phi \overset{\Gamma}{\nabla}{}_{\nu} \phi \\ &\quad + F_2(\phi, X) + F_3(\phi, X) \mathcal{L}_3^{\text{gal}\Gamma} + F_4(\phi, X) \mathcal{L}_4^{\text{gal}\Gamma}\end{aligned}$$

□ Kinetic coupling to Ricci scalar

$$\mathcal{L}(\overset{\Gamma}{R}, X) = f(X) \overset{\Gamma}{R} + P(X) \Leftrightarrow \mathcal{L} = f(X) R(g) + P(X) + \frac{6f_X^2}{f} \phi^\alpha \phi^\beta \phi_{\alpha\gamma} \phi_\beta^\gamma$$

Generalized k-essence? = Simplest theory of DHOST?

$$X := g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi$$

$$\phi_\mu = \nabla_\mu \phi, \quad \phi_{\mu\nu} = \nabla_\mu \nabla_\nu \phi$$

The counter term is **NOT artifact! Automatically obtained.**

One of the simplest DHOST without changing the speed of GWs.

Summary and Discussions

- Metric-affine formalism: metric and connection are independent.
- The covariant Galileon is unique due to the projective invariance.

- Class N-1/1a DHOST is

$$\mathcal{L}(g, \Gamma, \phi) = f_1(\phi, X) \overset{\Gamma}{R} + f_2(\phi, X) \overset{\Gamma}{G}{}^{\mu\nu} \overset{\Gamma}{\nabla}_{\mu} \phi \overset{\Gamma}{\nabla}_{\nu} \phi \\ + F_2(\phi, X) + F_3(\phi, X) \mathcal{L}_3^{\text{gal}\Gamma} + F_4(\phi, X) \mathcal{L}_4^{\text{gal}\Gamma}$$

- These theories may predict fermion-scalar coupling.

$$\mathcal{L}_{\text{int}} = \frac{i}{2} \kappa^{[\alpha\beta]}{}_{\beta}(g, \phi) j_{\alpha}$$

- Phenomenology?

$$j_{\alpha} = \bar{\psi} \gamma_{\alpha} \psi$$

Inflation or dark energy/matter in specific models?

Non-minimal fermion-scalar coupling?