Searching for general relativistic signatures on large scales

Jinn-Ouk Gong

APCTP, Pohang 790-784, Korea

CTPU IBS 20th August, 2015

590

Based on collaborations with S. G. Biern, J.-c. Hwang, D. Jeong and H. Noh

Introduction 000	Newtonian theory O	Relativistic theory	Relativistic theory with dark energy	Geodesic approach O	Conclusions O
Outlin	e				

- Introduction
- 2 Newtonian theory
- 3 Relativistic theory
 - Setup
 - Comoving gauge
 - Synchronous gauge
- 4 Relativistic theory with dark energy
 - Effects of dark energy
 - Non-linear power spectrum with dark energy
- 6 Geodesic approach
- 6 Conclusions

→ < ∃ →</p>

< 17 ▶



Planned galaxy surveys: DESI, HETDEX, LSST, Euclid, WFIRST...



Larger and larger volumes, eventually accessing the scales comparable to the horizon: beyond Newtonian gravity, fully general relativistic approach (or any modification) is necessary



Why non-linearity and gauge in LSS?

- Non-linearity is prominent in large scale structure thus accurate modeling of of non-linearity is very important
- GR is a gauge theory, thus observational quantities only make sense after choosing the coordinate systems

On large scales where non-linearity can be probed by observations with improved accuracy, density contrast $\delta \equiv (\rho - \bar{\rho})/\bar{\rho}$ deviates the Newtonian prediction

イロト イポト イヨト イヨト

 Introduction
 Newtonian theory
 Relativistic theory
 Relativistic theory with dark energy
 Geodesic approach
 Conclusions

 000
 0
 00000000
 00000000
 0
 0
 0

Why dark energy in non-linear regime?

- DE was negligible at very early times
- DE becomes significant at later stage when non-linearities in cosmic structure are developed

Naturally DE affects the evolution of gravitational instability, so that its effects emerge more prominently at non-linear level

Thus of our interest are:

- In the second second
- auge issue in GR
- effects of DE in non-linear regime of LSS

< ロ > < 同 > < 回 > < 回 > < 回 > <



3 basic equations for density perturbation $\delta \equiv \delta \rho / \bar{\rho}$, peculiar velocity *u* and gravitational potential Φ with a *pressureless* fluid

$$\dot{\delta} + \frac{1}{a} \nabla \cdot \boldsymbol{u} = -\frac{1}{a} \nabla \cdot (\delta \boldsymbol{u}) \qquad \text{continuity eq}$$
$$\dot{\boldsymbol{u}} + H\boldsymbol{u} + \frac{1}{a} \nabla \Phi = -\frac{1}{a} (\boldsymbol{u} \cdot \nabla) \boldsymbol{u} \qquad \text{Euler eq}$$
$$\frac{\Delta}{a^2} \Phi = 4\pi G \bar{\rho} \delta \qquad \text{Poisson eq}$$

Newtonian system is closed at 2nd order

$$\ddot{\delta} + 2H\dot{\delta} - 4\pi G\bar{\rho}\delta = -\frac{1}{a^2}\frac{d}{dt}[a\nabla\cdot(\delta \mathbf{u})] + \frac{1}{a^2}\nabla\cdot(\mathbf{u}\cdot\nabla\mathbf{u})$$

 \longrightarrow at linear order, $\delta_+ \propto a$ (growing) and $\delta_- \propto a^{-3/2}$ (decaying)

(Bernardeau et al. 2002)

Newtonian theory Relativistic theory Relativistic theory with dark energy •000000000

Basic non-linear equations

Based on the ADM metric

$$ds^{2} = -N^{2}(dx^{0})^{2} + \gamma_{ij} \left(N^{i} dx^{0} + dx^{i} \right) \left(N^{j} dx^{0} + dx^{j} \right)$$

the fully non-linear equations are (Bardeen 1980)

$$\begin{split} R - \overline{K}^{i}{}_{j}\overline{K}^{j}{}_{i} + \frac{2}{3}K^{2} - 16\pi GE &= 0 \\ \overline{K}^{j}{}_{i;j} - \frac{2}{3}K_{,i} &= 8\pi GJ_{i} \\ \\ \frac{K_{,0}}{N} - \frac{K_{,i}N^{i}}{N} + \frac{N^{;i}{}_{;i}}{N} - \overline{K}^{i}{}_{j}\overline{K}^{j}{}_{i} - \frac{1}{3}K^{2} - 4\pi G(E + S) &= 0 \\ \\ \frac{\overline{K}^{i}{}_{j,0}}{N} - \frac{\overline{K}^{i}{}_{j;k}N^{k}}{N} + \frac{\overline{K}_{jk}N^{i;k}}{N} - \frac{\overline{K}^{i}{}_{k}N^{k}{}_{;j}}{N} &= K\overline{K}^{i}{}_{j} - \frac{1}{N} \left(N^{;i}{}_{;j} - \frac{\delta^{i}{}_{j}}{3}N^{;k}{}_{;k} \right) + \overline{R}^{i}{}_{j} - 8\pi G\overline{S}^{i}{}_{j} \\ \\ \\ \frac{E_{,0}}{N} - \frac{E_{,i}N^{i}}{N} - K\left(E + \frac{S}{3}\right) - \overline{K}^{i}{}_{j}\overline{S}^{j}{}_{i} + \frac{\left(N^{2}J^{i}\right)}{N^{2}} &= 0 \\ \\ \\ \frac{J_{i,0}}{N} - \frac{J_{i;j}N^{j}}{N} - \frac{J_{j}N^{j}{}_{;i}}{N} - KJ_{i} + \frac{EN_{,i}}{N} + S^{j}{}_{i;j} + \frac{S^{j}{}_{i}N, j}{N} &= 0 \end{split}$$

Fluid quantities: $E \equiv n_{\mu} n_{\nu} T^{\mu\nu}$, $J_i \equiv -n_{\mu} T^{\mu}_i$, $S_{ii} \equiv T_{ii}$

Searching for general relativistic signatures on large scales

nar

・ロト ・回 ト ・ヨト ・ヨト

 Introduction
 Newtonian theory
 Relativistic theory
 Relativistic theory with dark energy
 Geodesic approach
 Conclusions

 000
 0
 00000000
 00000000
 0
 0
 0

Setup and perturbation variables

We consider scalar metric pert in Einstein-de Sitter universe

$$N = 1 + \alpha, \quad N_i = a^2 \beta_{,i}, \quad \gamma_{ij} = a^2 \left[(1 + 2\varphi) \delta_{ij} + \gamma_{,ij} \right]$$

The dynamical equations to be solved are:

Energy conservation $eq \rightarrow Continuity eq$ Trace of the Einstein $eq \rightarrow Euler eq$

We identify the perturbation variables as

$$\delta \equiv \frac{\rho - \bar{\rho}}{\bar{\rho}} \quad \text{with} \quad \rho \equiv -T_0^0$$
$$\theta \equiv \frac{\nabla \cdot \boldsymbol{u}}{a} = -3H - K$$

< ロ > < 同 > < 回 > < 回 > < 回 > <

Strategy for non-linear perturbations

Relativistic theory

000000000

Newtonian theory

With the linear solution the same as the standard one

$$\delta_1(\mathbf{k}, a) = D_1(a)\delta_1(\mathbf{k}, a_0)$$

Relativistic theory with dark energy

we expand $\delta = \delta_1 + \delta_2 + \cdots$ using symmetric kernels

$$\delta(\mathbf{k}, a) = \sum_{n=1}^{\infty} D^n(a) \int \frac{d^3 d_1 \cdots d^3 q_n}{(2\pi)^{3(n-1)}} \delta^{(3)} \left(\mathbf{k} - \mathbf{q}_1 - \cdots \mathbf{q}_n\right) \\ \times F_n(\mathbf{q}_1, \cdots \mathbf{q}_n) \delta_1(\mathbf{q}_1) \cdots \delta_1(\mathbf{q}_n)$$

Then correlation functions are

$$\langle \delta(\mathbf{k})\delta(\mathbf{k}') \rangle = (2\pi)^3 \delta^{(3)}(\mathbf{k} + \mathbf{k}')P(\mathbf{k}) \text{ with } P = P_{11} + \underbrace{P_{22} + P_{13}}_{1-\text{loop}} + \cdots$$

$$\langle \delta(\mathbf{k}_1) \delta(\mathbf{k}_2) \delta(\mathbf{k}_3) \rangle = (2\pi)^3 \delta^{(3)}(\mathbf{k}_{123}) B(k_1, k_2, k_3)$$

with $B = B_{112} + \underbrace{B_{222} + B_{123} + B_{114}}_{1-\text{loop}} + \cdots$

Introduction 000	Newtonian theory O	Relativistic theory	Relativistic theory with dark energy	Geodesic approach O	Conclusions O	
Comoving gauge						

We set the gauge condition as

$$\gamma = 0$$
 and $T_i^0 = 0$

Kernels are found to be (Jeong, JG, Noh & Hwang 2011, Biern, JG & Jeong 2014)

$$F_{2} = \frac{5}{7} + \frac{q_{1} \cdot q_{2}}{2q_{1}q_{2}} \left(\frac{q_{1}}{q_{2}} + \frac{q_{2}}{q_{1}}\right) + \frac{2}{7} \left(\frac{q_{1} \cdot q_{2}}{q_{1}q_{2}}\right)^{2}$$

$$F_{3} = F_{3N} + F_{3GR} \text{ where } F_{3GR} \propto k_{H}^{2} \text{ with } k_{H} \equiv aH$$

$$F_{4} = F_{4N} + (\cdots)k_{H}^{2} + (\cdots)k_{H}^{4}$$

- Those w/o φ are identical to the Newtonian kernels
- Newtonian kernels are the same as those found in the standard perturbation theory based on the Newtonian gravity
- GR contributions appear from 3rd order, prop to $k_H \equiv aH$

・ロット語 マイヨット ヨット シック



Power spectrum with leading corrections in CG



(Jeong, JG, Noh & Hwang 2011)

∃ → < ∃</p>

5990





Searching for general relativistic signatures on large scales

Leading bispectrum in various gauges



Introduction 000	Newtonian theory O	Relativistic theory	Relativistic theory with dark energy	Geodesic approach O	Conclusions O
Synch	ronous ga	uge			

We set the gauge condition as

$$g_{00} = -1$$
 and $g_{0i} = 0$

Kernels are found to be (Hwang, Noh, Jeong, JG & Biern 2014)

$$F_{2} = \frac{5}{7} + \frac{2}{7} \frac{(\boldsymbol{q}_{1} \cdot \boldsymbol{q}_{2})^{2}}{q_{1}^{2} q_{2}^{2}}$$

$$F_{3} = F_{3N} + F_{3GR,\varphi} + F_{3GR,no\varphi}$$

- Newtonian kernels are *different* from standard ones
- Some GR contributions are not from φ but from non-linear coupling w/o k_H (thus time independent)

<ロト < 同ト < 巨ト < 巨ト -



Power spectrum with leading corrections in SG



(Hwang, Noh, Jeong, JG & Biern 2014)

Jinn-Ouk Gong

Introduction Newtonian theory Relativistic theory Relativistic theory with dark energy Geodesic approach Conclusions 000 0 00000000 00000000 0 0 0

Newtonian interpretation of CG and SG

The problem lies in the Newtonian contributions

$$\dot{\delta} + \frac{1}{a}(1+\delta)\nabla \cdot \boldsymbol{u} = 0, \quad \dot{\theta} + 2H\theta + 4\pi G\bar{\rho}\delta + \frac{1}{a^2}\boldsymbol{u}^{i,j}\boldsymbol{u}_{j,i} = (\text{NL terms})$$



(Hwang, Noh, Jeong, JG & Biern 2014)

もしゃ 山下 エルト 山下 キャット

Introduction Newtonian theory o Relativistic theory o Relativistic theory with dark energy o Conclusion o Conclusion o Newtonian interpretation of CC and SC

Newtonian interpretation of CG and SG

The problem lies in the Newtonian contributions

$$\dot{\delta} + \frac{1}{a}(1+\delta)\nabla \cdot \boldsymbol{u} = 0, \quad \dot{\theta} + 2H\theta + 4\pi G\bar{\rho}\delta + \frac{1}{a^2}\boldsymbol{u}^{i,j}\boldsymbol{u}_{j,i} = (\text{NL terms})$$

$$\Downarrow \quad \frac{d}{dt} \rightarrow \frac{d}{dt} + \frac{1}{a}\boldsymbol{u}\cdot\nabla \quad \text{transformation to convective derivative}$$

$$\dot{\delta} + \frac{1}{a}\nabla \cdot [(1+\delta)\boldsymbol{u}] = 0, \quad \dot{\theta} + 2H\theta + 4\pi G\bar{\rho}\delta + \frac{1}{a^2}\nabla \cdot [(\boldsymbol{u}\cdot\nabla)\boldsymbol{u}] = (\text{NL terms})$$



(Hwang, Noh, Jeong, JG & Biern 2014)

Jinn-Ouk Gong

DQC

Introduction Newtonian theory Relativistic theory with dark energy Geodesic approach Ocnclusions Ococococo Putting dark energy on the table

Previous strategy is not complete

- ACDM power spectrum in EdS background
- Matter domination all the way

But we know the universe has been dominated by DE for a long time

$$\rho = \rho_m \longrightarrow \rho = \rho_m + \rho_{de}$$
 with $p_{de} = w \rho_{de}$

For simplicity

- No DE perturbation: $\rho_{dm} = \bar{\rho}_{de}$ (cf. Park, Hwang, Lee & Noh 2009)
- **2** Comoving gauge: $T^0_i = 0$

イロト イポト イヨト イヨト

Introduction Newtonian theory Relativistic theory Relativistic theory with dark energy Geodesic approach Conclusions 000 0 000000000 0

Dark energy changes the game

DE provides different BG from both EdS and ACDM:

$$\mathcal{H}^2 = \frac{8\pi G}{3} a^2 \left(\bar{\rho}_m + \bar{\rho}_{de} \right) \quad \text{and} \quad \mathcal{H}' = -\frac{1}{2} \mathcal{H}^2 (1 + 3w)$$

DE permeates all order in perturbation: e.g. energy conservation

$$\delta' - \kappa (1 - \lambda) = (\text{non-linear terms}) \text{ where } \lambda \equiv (1 + w) \left(1 - \frac{1}{\Omega_m}\right)$$

Thus away from EdS ($\Omega_m = 1$) and Λ CDM (w = -1) the effects of general, dynamical DE are *manifest*: we use the parametrization

(Chevallier & Polarski 2001, Linder 2003)

$$w(a) = w_0 + (1 - a)w_a$$

イロト イポト イヨト イヨト

 Introduction
 Newtonian theory
 Relativistic theory
 Relativistic theory with dark energy
 Geodesic approach
 Conclusions

 000
 0
 000000000
 00000000
 0
 0
 0

Non-linear solutions with DE

Curvature perturbation is not conserved: from energy constraint

$$\varphi = -\frac{\mathcal{H}^2 f}{1-\lambda} \left[1 + \frac{3}{2} (1-\lambda) \frac{\Omega_m}{f} \right] \Delta^{-1} \delta \neq \text{constant}$$

Thus δ receives a) curvature evolution effects from 3rd order and b) general, dynamical DE effects from BG and linear order:

$$\delta'' + \left(\mathcal{H} + \frac{\lambda'}{1-\lambda}\right)\delta' - \frac{3}{2}(1-\lambda)\mathcal{H}^2\Omega_m\delta = \underbrace{\mathcal{N}_N + \mathcal{N}_{\varphi} + \mathcal{N}_{\varphi'} + \mathcal{N}_{\lambda}}_{=\text{non-linear source terms}}$$

	Newtonian	EdS	ΛCDM	DE
\mathcal{N}_N	0	0	0	0
$\mathcal{N}_{oldsymbol{arphi}}$	Х	0	0	0
$\mathcal{N}_{oldsymbol{arphi}}$	Х	Х	Х	0
\mathcal{N}_{λ}	Х	Х	Х	0

< ロ > < 同 > < 回 > < 回 > < 回 > <

Introduction 000	Newtonian theory O	Relativistic theory	Relativistic theory with dark energy	Geodesic approach O	Conclusions O			
Relativistic kernels								

2nd and 3rd order solutions are (Biern & JG 2015)

$$\begin{split} \delta_{2}(\mathbf{k}, a) &= D_{1}^{2} \sum_{i=a}^{b} c_{2i}(a) \int \frac{d^{3}q_{1} d^{3}q_{2}}{(2\pi)^{3}} \delta^{(3)}(\mathbf{k} - \mathbf{q}_{12}) F_{2i}(\mathbf{q}_{1}, \mathbf{q}_{2}) \delta_{1}(\mathbf{q}_{1}) \delta_{1}(\mathbf{q}_{2}) \\ \delta_{3}(\mathbf{k}, a) &= D_{1}^{3} \sum_{i=a}^{f} c_{3i}(a) \int \left[\cdots F_{3i} \cdots 3 \delta_{1}' s \right] \qquad c_{ni} \equiv \frac{D_{ni}}{D_{1}^{n}} \\ &+ D_{1}^{3} \mathcal{H}^{2} \sum_{i=a}^{b} c_{3i}^{\varphi}(a) \int \left[\cdots F_{3i}^{\varphi} \cdots 3 \delta_{1}' s \right] \qquad c_{3i}^{\varphi} \equiv \frac{D_{3i}^{\varphi}}{D_{1}^{3} \mathcal{H}^{2}} \end{split}$$

In the EdS universe *c*'s are fixed as certain numbers ($c_{2a} = 3/7...$) and (also in Λ CDM) c_{ni} terms become purely Newtonian _[Kamionkowski & Buchalter 1999 (2nd) and Takahashi 2008 (3rd)] and only c_{3i}^{φ} terms remain relativistic

N. B. λ is completely entangled and cannot be separated like φ

・ロト ・ 同ト ・ ヨト ・ ヨト … ヨ

Relativistic theory Relativistic theory with dark energy 00000000

One-loop corrected power spectrum: versus ΛCDM



- Overall almost constant deviation on large scales $(k \leq 0.1 h/\text{Mpc})$
- Deviation becomes significant on $k \gtrsim 0.1 h/Mpc$, close to baryon acoustic oscillations
- $w_0 > -1 / w_a > 0$ ($w_0 < -1 / w_a < 0$) give smaller (larger) P(k)

nar

Introduction Newtonian theory Relativistic theory with dark energy Geodesic approach Conclusie One a locar comment of a comment of a comment of the commen

One-loop corrected power spectrum: versus EdS

In Newtonian studies, usually EdS power spectrum is transferred to an arb model by replacing $a \rightarrow D_1(a)$:

 $P(k, a) = D_1^2(a)P_{11}(k) + D_1^4(a)[P_{22}(k) + P_{13}(k)]_{\rm EdS}$



- For Λ CDM, only φ drives difference so almost identical to EdS
- For general DE, the difference notably increases from $k \approx 0.1 h/\text{Mpc}$



Observations are made i.t.o. redshift (Kaiser 1987, Heavens, Matarrese & Verde 1998

$$\delta_s = \delta_r - \partial_{\parallel} U + \text{higher order terms}$$

where $\delta_r = b\delta$, $U \equiv \frac{\hat{\boldsymbol{n}} \cdot \boldsymbol{v}}{\mathcal{H}}$ and $\partial_{\parallel} \equiv \hat{\boldsymbol{n}} \cdot \nabla$

Then the observable galaxy power spectrum in the redshift space

$$P_{s}(k,\mu,a) = P_{s11}(k,\mu,a) + P_{s22}(k,\mu,a) + P_{s13}(k,\mu,a)$$

with $\mu \equiv \hat{\boldsymbol{n}} \cdot \boldsymbol{k} / k$, thus no longer isotropic

- $\mu = 1$: line-of-sight direction, most dominant
- $\mu = 0$: perp to LoS

Thus the deviation from ACDM becomes larger for LoS spectrum

< ロ > < 同 > < 回 > < 回 > .



Deviation is enhanced as large as 10% at around BAO scales

$w_a = 0$ and varying w_0			$w_0 = -1$ and varying w_a			
<i>k</i> [<i>h</i> /Mpc]	$w_0 = -1.2$	$w_0 = -0.8$	<i>k</i> [<i>h</i> /Mpc]	$w_a = -1.0$	$w_a = -0.5$	$w_a = 0.5$
0.1	6.8%	-10.2%	0.1	9.5%	5.8%	-11.5%
0.2	11.6%	-15.0%	0.2	14.9%	8.8%	-15.3%
0.3	16.0%	-19.4%	0.3	20.1%	11.6%	-19.0%

Searching for general relativistic signatures on large scales

イロト イポト イヨト イヨ

5990

Introduction Newtonian theory of Relativistic theory with dark energy occosed approach of Conclusions of Conclu

Observable galaxy number density

We observe as if photons come to us along a straight, unperturbed geodesic...



▶ ∢ ≣

Introduction Newtonian theory oo
Relativistic theory with dark energy ooooooo
Geodesic approach oo
Conclusions o

Observable galaxy number density

We observe as if photons come to us along a straight, unperturbed geodesic... but in fact the path is distorted due to perturbations at the locations of the observer and the source, and in between

(Yoo et al. 2009, Bonvin & Durrer 2011, Bertacca, Maartens & Clarkson 2014, Yoo & Zaldarriaga 2014...)



• • • • • • • • • • • • • •

Introduction Newtonian theory of Relativistic theory with dark energy occosed approach of Conclusions of Conclu

Observable galaxy number density

We observe as if photons come to us along a straight, unperturbed geodesic... but in fact the path is distorted due to perturbations at the locations of the observer and the source, and in between

(Yoo et al. 2009, Bonvin & Durrer 2011, Bertacca, Maartens & Clarkson 2014, Yoo & Zaldarriaga 2014...)



See S. G. Biern's presentation for detail

Introduction 000	Newtonian theory O	Relativistic theory	Relativistic theory with dark energy	Geodesic approach O	Conclusions •
Conclu	usions				

- As galaxy surveys become deeper and deeper, fully GR description is relevant
- Gauge dependence at non-linear order:
 - In CG the standard perturbation theory is reproduced
 - Pure GR corrections are heavily suppressed in almost all cases
 - Naively using SG leads to pathologies
 - Transformation by hands cures the problem
- With general dark energy:
 - Dark energy background greatly affects GR contributions
 - Notable difference of a few percent near BAO scales
 - Detectable signatures of judging Λ or not
- Geodesic approach based on observable quantities should help

イロト イポト イヨト イヨト 二日