

Dark Matter in the Type Ib Seesaw Model



Bowen Fu

付博文


Based on M. Chianese, B. Fu, and S. F. King, (2021), arXiv:2102.07780 [hep-ph]

The 26th International Symposium on Particle Physics, String Theory, and Cosmology (PASCOS 2021)

16 June 2021

Dark Matter Production

- ❖ Freeze-out: Thermal equilibrium
- ❖ Freeze-in: Never in thermal equilibrium




- ❖ The observed DM relic abundance gives a constraint to DM yield




$$\Omega_{\text{DM}}^{\text{obs}} h^2 = 0.120 \pm 0.001$$

Neutrino Portal

- ❖ General neutrino portal
- ❖ heavy scalar scenario
- ❖ Dark matter producing process




ν -Yukawa process $\propto Y^2 y^2$



dark sector process $\propto y^4$

Why Type Ib Seesaw Model?


- ❖ Traditional type I seesaw mechanism (type Ia)



$$m \propto \frac{Y^2 v^2}{M_N}$$

- ❖ $m \sim 0.1 \text{ eV}, M_N \sim 1 \text{ GeV} \rightarrow Y \sim 10^{-7}$
- ❖ $M_N > 4 \text{ TeV}$ for ν -Yukawa dominance

- ❖ Type Ib seesaw mechanism



$$m \propto \frac{Y_1 Y_2 v_1 v_2}{M_N}$$

- ❖ One of Y_1, Y_2 can be small while the other one is large

Type Ib Seesaw Model with a Neutrino Portal

❖ Particles and symmetries

	Q_α	$u_{R\beta}$	$d_{R\beta}$	L_α	$e_{R\beta}$	Φ_1	Φ_2	N_{R1}	N_{R2}	ϕ	$\chi_{L,R}$
$SU(2)_L$	2	1	1	2	1	2	2	1	1	1	1
$U(1)_Y$	$\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{3}$	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	$-\frac{1}{2}$	0	0	0	0
Z_3	1	ω	ω	1	ω	ω	ω^2	ω^2	ω	ω	ω^2
Z_2	+	+	+	+	+	+	+	+	+	−	−

❖ Seesaw Lagrangian and neutrino portal

$$\mathcal{L}_{\text{seesaw Ib}} = -Y_{1\alpha} \bar{L}_\alpha \Phi_1 N_{R1} - Y_{2\alpha} \bar{L}_\alpha \Phi_2 N_{R2} - M_N \overline{N_{R1}^c} N_{R2} + \text{h.c.}$$

$$\mathcal{L}_{N_R \text{portal}} = y_1 \phi \overline{\chi_R} N_{R1}^c + y_2 \phi \overline{\chi_L} N_{R2} + \text{h.c.}$$

❖ The two RH neutrinos form a Dirac pair $\mathcal{N} = (N_{R1}^c, N_{R2})$

$$\mathcal{L}_{\text{seesaw Ib}} = -Y_{1\alpha}^* \bar{L}_\alpha^c \Phi_1^* \mathcal{N}_L - Y_{2\alpha} \bar{L}_\alpha \Phi_2 \mathcal{N}_R - M_N \overline{\mathcal{N}_L} \mathcal{N}_R + \text{h.c.}$$

$$\mathcal{L}_{N_R \text{portal}} = y \phi \overline{\chi} \mathcal{N} + \text{h.c.}$$

Constraints


❖ Oscillation data

$$Y_{1\alpha} \rightarrow Y_1$$

$$Y_{2\alpha} \rightarrow Y_2$$

$$\frac{Y_1 Y_2 v_1 v_2}{M_N} = 3.0 \times 10^{-11} \text{ GeV}$$

❖ Muon decay



$$|\eta_{e\mu}| = \frac{|Y_{1e} Y_{1\mu}^* v_1^2 + Y_{2e} Y_{2\mu}^* v_2^2|}{4M_N^2} \lesssim 8.4 \times 10^{-6}$$

❖ Collider data on neutrino mixing


$$U_\alpha^2 = \sum_{i=L,R} |U_{\alpha i}|^2, \quad \alpha = e, \mu, \tau$$

$$U_\alpha^2 = (a_\alpha + b_\alpha \cos \delta_M) \frac{v_1^2 Y_1^2}{M_N^2} + (a_\alpha - b_\alpha \cos \delta_M) \frac{v_2^2 Y_2^2}{M_N^2}$$

$$M_{\text{diag}}^\nu = U^T M^\nu U$$

$$U_\alpha^2 \geq \sqrt{a'_\alpha - b'_\alpha \cos \delta_M} \frac{1 \text{ GeV}}{M_N}$$

Required Portal Coupling



❖ $\tan\beta = v_2/v_1 = 10$

❖ If $m_\phi \gg m_\chi, M_N$, then the dark matter production only depends on m_ϕ/m_χ instead of m_ϕ or m_χ


- dark sector dominance $y^4 \simeq 3.9 \times 10^{-23} \frac{m_\phi}{m_\chi}$
- ν -Yukawa dominance $(Y_1^2 + Y_2^2)^2 \simeq 5.5 \times 10^{-22} \frac{m_\phi}{m_\chi}$

❖ M_N is 1, 10^2 , 10^4 GeV

❖ For heavier M_N , the ν -Yukawa dominance is favoured in most of the parameter space


$$\frac{m_\phi}{m_\chi} \gtrsim 6.9 \times 10^{-9} \left(\frac{M_N}{1\text{GeV}} \right)^2 \frac{1}{\sin^2 2\beta}$$

Relation to Experiments



- ❖ ν -Yukawa dominance is allowed above the coloured dashed lines
- ❖ Less constrained as $\tan\beta$ increases
- ❖ The strongest constraint is given by ν_μ mixing

Relation to Experiments



- ❖ ν -Yukawa dominance is allowed above the coloured dashed lines
- ❖ More constrained as m_ϕ/m_χ increases
- ❖ The strongest constraint is given by ν_μ mixing

Summary

- ❖ We discussed dark matter production in a minimal extension of type Ib seesaw model
- ❖ The DM- ν relation is constrained by experimental results when the heavy neutrino is GeV scale
- ❖ The results can be further constrained by future experiments
- ❖ The type Ib seesaw mechanism can be realised effectively to produce leptogenesis

Thank You!