EMIS 2022 at RAON



Contribution ID: 101

Type: Invited

Studies of the two-step scheme with a ¹³²Sn beam for next-generation RI-beam production method in the medium-heavy very-neutron-rich region

Tuesday, 4 October 2022 09:00 (30 minutes)

The usefulness of the two-step scheme with a ¹³²Sn beam was investigated [1], which was proposed for efficient production of medium-heavy very-neutron-rich radioactive isotopes (RI) [2] as an alternative method to the direct production by means of in-flight fission of a ²³⁸U beam (one-step scheme). The system of the two-step scheme consists of an isotope-separation online (ISOL) system and an in-flight fragment separator. Long-lived neutron-rich RIs (e.g., ¹³²Sn) are produced by ISOL with a thick U target and a high-intensity proton beam in the first step, and more neutron-rich RI beams (e.g., ¹²⁸Pd) are produced by a projectile fragmentation from the re-accelerated less-exotic RI beams in the second step.

We measured production cross sections of very neutron-rich RIs around a N = 82 region beyond ¹²⁵Pd, up to which the cross sections had already been measured at GSI [3], with a 278-MeV/nucleon ¹³²Sn beam produced by the BigRIPS separator [4] impinging on a 5.97-mm Be target. The yields obtained by the two-step and one-step schemes were estimated based on the measured cross sections, and we examined whether and to what extent the two-step scheme at future 1-MW beam facilities can reach further into the neutron-rich regions. This comparison suggests that the two-step scheme with the ¹³²Sn beam provides yields 40-times higher than those with the one-step scheme for the very neutron-rich N = 82 region. Moreover, by using various RI beams over the nuclear chart from ISOL, certain regions of very neutron-rich RIs around N = 50, 60, 82, and 90 regions, including the supernova *r*-process path, can be produced with greater yields than by the one-step approach.

References

- [1] H. Suzuki et al., Phys. Rev. C 102, 064615 (2020).
- [2] K. Helariutta et al., Eur. Phys. J. A 17, 181 (2003).
- [3] D. Pérez-Loureiro et al., Phys. Lett. B 703, 552 (2011).
- [4] T. Kubo, Nucl. Instr. and Meth. B 204, 97 (2003).

Primary authors: SUZUKI, Hiroshi (RIKEN Nishina Center); BAZIN, Daniel (FRIB / MSU); FUKUDA, Naoki (RIKEN); Prof. HENNING, Walter, F. (Argonne National Laboratory, Technische Universitaet Muenchen); IMAI, Nobuaki (CNS, the University of Tokyo); Mr INABE, Naohito (RIKEN Nishina Center); Dr KAWATA, Keita (CNS, University of Tokyo); Dr KITAMURA, Noritaka (CNS, University of Tokyo); Dr KOMATSUBARA, Tetsuro (RIKEN Nishina Center); KORKULU, Zeren (IBS, CENS); Dr KUBO, Toshiyuki (RIKEN Nishina Center); Dr KUSAKA, Kensuke (RIKEN Nishina Center); MICHIMASA, Shin'ichiro (The University of Tokyo CNS); Prof. NOLEN, Jerry (Argonne National Laboratory); Mr OHTAKE, Makoto (RIKEN Nishina Center); Dr SATO, Hiromi (RIKEN Nishina Center); SHIMIZU, Yohei (RIKEN); SUMIKAMA, Toshiyuki (RIKEN); TAKEDA, Hiroyuki (RIKEN); TARASOV, Oleg (FRIB / MSU); UENO, Hideki (RIKEN Nishina Center); Dr YANAGISAWA, Yoshiyuki; YOSHIDA, Koichi (RIKEN); AHN, Deuk Soon (Center for Exotic Nuclear Studies, IBS)

Presenter: SUZUKI, Hiroshi (RIKEN Nishina Center)

Session Classification: Session 5