

GRAVITATIONAL WAVE SIGNATURES OF REHEATING


Based on arXiv:2305.09712
with Manuel A. Buen-Abad and Anson Hook

Jae Hyeok Chang


Johns Hopkins University and University of Maryland

2023/06/13 PPC 2023


Chronology of the Universe



Chronology of the Universe




Chronology of the Universe




Chronology of the Universe

- We have observational data only after BBN




Chronology of the Universe

- We have observational data only after BBN




Chronology of the Universe

- We have observational data only after BBN
- How do we probe earlier Universe?




Chronology of the Universe

- We have observational data only after BBN
- How do we probe earlier Universe?
- We need a messenger from earlier Universe




Gravitational Waves!

- Universe is transparent to GW for all time
- We can probe earlier time if GW are produced from some early-time events




Gravitational Waves!


- Universe is transparent to GW for all time
- We can probe earlier time if GW are produced from some early-time events
- e.g. 1st order phase transitions



GW from 1st order PT



GW from 1st order PT




- GW are produced from bubble dynamics (bubble collisions, sound waves, and turbulences)

If we have a 1st order PT at T_c ,
it must occur twice:

If we have a 1st order PT at T_c ,
it must occur twice:
once during the reheating period
and once again during
the subsequent cooling period


Temperature history during reheating




Temperature history during reheating

$$\dot{\rho}_\chi + 3H\rho_\chi = -\Gamma_\chi\rho_\chi$$


$$\dot{\rho}_r + 4H\rho_r = \Gamma_\chi\rho_\chi$$




Temperature history during reheating




Temperature history during reheating




Temperature history during reheating



1st order PT during the reheating



GW from PT during the reheating





- GW are produced from the similar ways, but there're several differences

Heating PT and Cooling PT

- To distinguish, we define
 - **hPT** as phase transition during the reheating
 - **hGW** as gravitational waves from **hPT**
 - **cPT** as Phase transition during the cooling
 - **cGW** as gravitational waves from **cPT**


Difference between hGW and cGW

- **hGW** redshift more than **cGW**
 - **hGW** are produced earlier, and GW redshift $\propto a^{-4}$
 - This is why **hGW** are not considered normally
 - We can reduce this redshift factor by having $T_c \lesssim T_{\max}$
 - Yet, we need something else to avoid fine-tuning



Difference between hGW and cGW

- Different bubble dynamics
 - The bubble wall in cPT feels friction, while the wall in hPT feels anti-friction
 - Thus, normally the wall speed in cPT ($v_{w,cPT}$) saturates at a constant value, while $v_{w,hPT}$ enters a runaway regime
 - The bubble wall in hPT gets energy from the radiation, which eventually converts to GW



Toy Model for PT

$$V(\Phi) = \frac{\mu^2}{2} (T - T_0)^2 \Phi^2 + \frac{A}{3} T \Phi^3 + \frac{\lambda}{4!} \Phi^4$$

- Φ : A scalar in the radiation r that drives PT
- $\mu^2 = \frac{1}{12} \sum_i c_i N_i y_i^2$, $A = \frac{1}{4\pi} \sum_B N_B y_B^3$
- We define $\Delta \equiv \frac{4A^2}{3\lambda\mu^2}$, which controls physics of PT
- T_0 : Binodal temperature
- $T_c = T_0 / \sqrt{1 - \Delta}$: Critical temperature
- $T_1 = T_0 \sqrt{8 / (8 - 9\Delta)}$: Spinodal temperature

Free Parameters


$$T_c, \quad \Gamma_\chi, \quad T_c/T_{\max}, \quad \{\mu, A, \lambda\}, \quad g_*, \quad \nu_{w,\text{cPT}}$$

- We choose
 - $T_c \sim 1 \text{ TeV}$
 - $\Gamma_\chi \sim H_i$
 - $\mu = \lambda = 1$
 - $g_* = 10$
 - $\nu_{w,\text{cPT}} = 0.05$
- Vary T_c/T_{\max} and $\Delta \equiv \frac{4A^2}{3\lambda\mu^2}$


Gravitation Waves from PT

- The dominant contribution for **hGW** comes from bubble collisions as it's in the runaway regime
- The bubble wall for **cPT** has a constant wall velocity, thus sound waves dominates for **cGW**
- We calculate SNR in BBO by using Ω_{PIS}



Gravitation Waves from PT





Results





Results: different T_c



Results: different Γ_χ/H_i



Results: varying Γ_χ/H_i for fixed Δ



Conclusions

- 1st order phase transition may occur twice during the reheating and during the cooling
- If we can measure GW from both PTs, we can learn physics of reheating
- In this work, we study the properties of GW production during the reheating

THANK YOU

BACK UP

Toy Model for Reheating

- A non-relativistic reheaton χ decays to a radiation r , which has 1st order phase transition

$$\dot{\rho}_{\chi} + 3H\rho_{\chi} = -\Gamma_{\chi}\rho_{\chi}$$

$$\dot{\rho}_r + 4H\rho_r = \Gamma_{\chi}\rho_{\chi}$$

- Γ_{χ} is the reheaton decay rate
- Radiation r thermalizes with SM sector later
- χ can be an inflaton, but not necessarily


Bubble Nucleation Rate

$$\frac{\Gamma}{\mathcal{V}} \approx T^4 \left(\frac{S}{2\pi} \right)^{3/2} e^{-S}$$


$$h(t) = \exp \left[- \int_{t_c}^t dt' \frac{\Gamma}{\mathcal{V}}(t') \frac{4\pi}{3} v_w^3 (t - t')^3 \right]$$

- h is the fraction of the volume of the Universe found in the metastable phase
- Define t_{PT} at which $h(t_{PT}) = 1/e$

Bubble Nucleation Rate



Temperature history according to γ



Daisy, runaway, and κ_Φ

