


# Composite Dirac neutrinos with QCD axion

Tae Hyun Jung


KIAS

Ref S. Chakraborty, THJ, T. Okui, arXiv: 2108.04293


# Outline



# Outline




# Composite Dirac neutrinos




# Why are neutrinos so light?

$$y_\nu H l \nu^c + \text{h. c.} + M_\nu \nu^c \nu^c$$

1. See-saw mechanism (Majorana neutrinos):  $M_\nu \gg y_\nu v_h$



$$\Rightarrow m_\nu \sim y_\nu v_h \left( \frac{y_\nu v_h}{M_\nu} \right)$$



**$2.0 \neq y_\nu < 10^{-13}$  without  $M_\nu$**

N. Arkani-Hamed, Y. Grossman (1999)

**Composite  $\nu^c$**

# Composite Dirac neutrinos

N. Arkani-Hamed, Y. Grossman (1999)

$$\nu^c \propto \psi\psi\chi$$

$\psi, \chi$ : fundamental Weyl

$\nu^c$ : composite Weyl (like proton/neutron in QCD:  $p \propto uud$ ,  $n \propto udd$ )



Effective description  
after confinement

$$H l \nu^c$$

$\leftrightarrow$

$$\frac{1}{M^3} H l (\psi\psi\chi) \longrightarrow \text{Dim: 7}$$

$$(\psi\psi\chi) \sim \Lambda_c^3 \nu^c$$



$$y_\nu \sim \left(\frac{\Lambda_c}{M}\right)^3$$

$$\frac{\Lambda_c}{M} \sim 10^{-4} \Rightarrow m_\nu \sim 0.1 \text{ eV}$$

\* Majorana operator  $(\psi\psi\chi)^2$  : dim 9

**Q: How can those composite fermions remain massless?**

cf) baryons in QCD

 **A special setup required**

# Composite Dirac neutrinos

Arkani-Hamed, Grossman (1999)

	Gauge
	$SU(6)$
$\psi$	$\bar{\mathbf{6}} \times 2$ (Two generation)
$\chi$	$\mathbf{15}$
$\nu^c \propto \psi\psi\chi$	$\mathbf{1} \times 3$ (Three generation)

# Composite Dirac neutrinos

Arkani-Hamed, Grossman (1999)

	Gauge	Global	
	$SU(6)$	$U(1)_X$	
$\psi$	$\bar{\mathbf{6}}$	$-2/3$	$\times 2$ (Two generation)
$\chi$	$\mathbf{15}$	$1/3$	
$\nu^c \propto \psi\psi\chi$	$\mathbf{1}$	$-1$	$\times 3$ (Three generation)


$$A(U(1)_X SU(6)^2) = 0$$

# Composite Dirac neutrinos

Arkani-Hamed, Grossman (1999)


	Gauge	Global	
	$SU(6)$	$U(1)_X$	
$\psi$	$\bar{\mathbf{6}}$	$-2/3$	$\times 2$ (Two generation)
$\chi$	$\mathbf{15}$	$1/3$	
$\nu^c \propto \psi\psi\chi$	$\mathbf{1}$	$-1$	$\times 3$ (Three generation)

$$A(U(1)_X SU(6)^2) = 0$$



**UV**

$$= -3$$



**IR**

't Hooft anomaly matching conditions

S. Dimopoulos, S. Raby, L Susskind (1980)

# Composite Dirac neutrinos

Arkani-Hamed, Grossman (1999)

	Gauge	Global	
	$SU(6)$	$U(1)_L$	
$\psi$	$\bar{\mathbf{6}}$	$-2/3$	$\times 2$ (Two generation)
$\chi$	$\mathbf{15}$	$1/3$	
$\nu^c \propto \psi\psi\chi$	$\mathbf{1}$	$-1$	$\times 3$ (Three generation)
$l$	$\mathbf{1}$	$+1$	$\times 3$ (Three generation)

Interaction with SM:  $\frac{\Lambda_c^3}{M^3} Hl \nu_c \sim \frac{1}{M^3} Hl (\psi\psi\chi)$

$B - L$  : good


$L_c, B - L_{SM}$  : broken explicitly

$(B - L_{SM})$ -genesis at  $T \sim M$  scale?


possible by itself, but requires some tuning in UV

Grossman, Tsai (2008)

# Kinetic misalignment



# Kinetic misalignment




Co, Harygaya, 1910.02080

Co, Harygaya, Hall, 1910.14152

Potential of PQ breaking field  $\Phi$

# Kinetic misalignment



randomly picked up  
by quantum fluctuation  
during inflation

$V(\Phi)$

$\text{Re}[\Phi]$


$\text{Im}[\Phi]$

Potential of PQ breaking field  $\Phi$

Co, Harygaya, 1910.02080

Co, Harygaya, Hall, 1910.14152

# Kinetic misalignment



randomly picked up  
by quantum fluctuation  
during inflation


Some explicit breaking  
e.g.  $\Phi^n / M_{Pl}^{n-4}$

Potential of PQ breaking field  $\Phi$

Co, Harygaya, 1910.02080

Co, Harygaya, Hall, 1910.14152

# Kinetic misalignment




Potential of PQ breaking field  $\Phi$

Co, Harygaya, 1910.02080

Co, Harygaya, Hall, 1910.14152

# Evolution of the axion motion




At  $T_i \ll f_a$ ,  $\dot{a}(T_i) = \dot{a}_i \neq 0$

PQ charge from the axion motion is approximately conserved, so

$$Y_{PQ} = \frac{n_{PQ}}{s} \simeq \frac{\dot{a}(T)f_a}{s} \simeq \text{const} \Rightarrow \dot{a}(T) \simeq \frac{Y_{PQ}}{f_a} s(T)$$

**free parameter in this talk**  
(fixed by initial condition)

Maybe some modification  
from QCD friction  
(Alexandros' talk)



# Axiogenesis


With  $\dot{a} \neq 0$  background,

Co, Harigaya (2019)

1. Chirality asymmetry is generated.

$$\frac{a}{f_a} G\tilde{G} \rightarrow \frac{a}{f_a} \partial_\mu J_A^\mu \rightarrow -\frac{\partial_\mu a}{f_a} J_A^\mu \rightarrow -\frac{\dot{a}}{f_a} (n_L - n_R)$$

2. Weak sphaleron converts only left-handed quarks into right-handed anti-leptons.




**3.  $(B + L)$ -genesis**

$$n_B \sim \mu_{B+L} T^2 \sim \frac{\dot{a}}{f_a} T^2 \sim Y_{PQ} \left(\frac{T}{f_a}\right)^2 s$$

$\dot{a} \approx Y_{PQ} \frac{s}{f_a}$

# Axion dark matter

$$\text{QCD axion: } m_a \sim \frac{m_\pi f_\pi}{f_a}$$




...

$$\eta_B = \frac{n_B}{s} \sim Y_{PQ} \left( \frac{T}{f_a} \right)^2$$


# Axion dark matter

QCD axion:  $m_a \sim \frac{m_\pi f_\pi}{f_a}$



# Axion dark matter

QCD axion:  $m_a \sim \frac{m_\pi f_\pi}{f_a}$




# Strong CP problem, baryon asymmetry, dark matter


$$m_a f_a \sim m_\pi f_\pi$$

$$Y_{PQ} \left( \frac{T_{EW}}{f_a} \right)^2 \sim 10^{-10}$$

$$m_a Y_{PQ} \sim 0.1 \text{ eV}$$




Co, Harygaya, 1910.02080  
Co, Harygaya, Hall, 1910.14152



## with composite Dirac $\nu$

$$L = L_{SM} + L_c$$



$$\frac{1}{M^3} H l (\psi \psi \chi)$$


$$L_{SM} \leftrightarrow L_c$$

**Active at high  $T > T_{LD} \sim 0.1M$**

LD: (composite) Lepton Decoupling (chemically)

# with composite Dirac $\nu$

$$L = L_{SM} + L_c$$




$$L_{SM} \leftrightarrow L_c$$

Active at high  $T > T_{LD} \sim 0.1M$

LD: (composite) Lepton Decoupling (chemically)

$$\frac{1}{M^3} H l (\psi \psi \chi)$$




- quarks
- anti-quarks
- anti-leptons
- composite sector

$$\begin{aligned}
 B &= 0, L = 0 \\
 L_{SM} &= 0, L_c = 0 \\
 B + L_{SM} &= 0 \\
 B - L_{SM} &= 0
 \end{aligned}$$


$$\begin{aligned}
 B &= -1, L = -1 \\
 L_{SM} &= 1, L_c = 0 \\
 B + L_{SM} &= -2 \\
 B - L_{SM} &= 0
 \end{aligned}$$

$$\begin{aligned}
 B &= -1, L = -1 \\
 L_{SM} &= 0, L_c = 1 \\
 B + L_{SM} &= -1 \\
 B - L_{SM} &= -1
 \end{aligned}$$


# Thermal History with composite Dirac $\nu$




# Baryogenesis with composite Dirac $\nu$



# Baryogenesis with composite Dirac $\nu$



# Baryogenesis with composite Dirac $\nu$




**$\Delta N_{\text{eff}}$  from  $\nu^c$  ?**



**Thermal production required  
for baryogenesis**

$$\Delta N_{\text{eff}}$$

If  $T_{kd} \gtrsim \Lambda_c$




$$\Delta N_{\text{eff}} \simeq 8.8$$

ruled out!

$$\Delta N_{\text{eff}}$$

If  $T_{kd} \ll \Lambda_c$




$$\Delta N_{\text{eff}} \gtrsim 0.14$$


Not ruled out,  
and testable in future

CMB stage IV:  $\Delta N_{\text{eff}} < 0.03$

# Summary



# Summary



$$m_a > \mu\text{eV}$$

$$f_a < 10^{12}\text{GeV}$$

**axiogenesis**

**&  $\Delta N_{\text{eff}} \gtrsim 0.14$**

**& No  $0\nu\beta\beta$**

**composite  $\nu^c$**