

2024 Workshop on (Mostly) Matroids

Monday, 19 August 2024 - Friday, 23 August 2024

IBS Science Culture Center

Book of Abstracts

Contents

Geometric methods for matroid theory	1
Is There Hope?	1
Characterising matroids representable over all fields of size at least four	1
On the matroid representation over finite rings	2
Towards Rota’s conjecture for gain-graphic matroids	2
Rigidity and reconstruction in matroids of highly connected graphs	2
Infinite matroids on lattices	3
Unbounded matroids, polymatroids and subspace arrangements	3
Tropical Ideals	4
Amalgams of matroids, fibre products and tropical graph correspondences	4
Speyer’s g conjecture via external activity of a pair of matroids	4
Symmetric tropical rank 2 matrices	5
Baker-Bowler theory for Lagrangian Grassmannians	5
Direct sum of q -matroids and linear sets	5
Almost all q -matroids are not representable	6
On a q -analogue of perfect matroid designs	6
The cyclic flats of L-polymatroids	6
Graphs embedded on surfaces and their delta-matroids	7
Graphs in surfaces, their one-face subgraphs, and the critical group	7
Clique minors in dense matroids	8
Chordal matroids arising from generalized parallel connections	8
Connectivity systems, path-width, and well-quasi-ordering	8
Average Hyperplane-Size in Complex-Representable Matroids	9

Convex geometry of building sets	9
Combinatorics of boundary algebras.	9
Complexity of log-concave inequalities in matroids	10
Chow rings and augmented Chow rings of matroids are equivariant γ -positive	10
Representations on the cohomology of the moduli space of pointed rational curves, permutation representations and log concavity	11
Intrinsic properties of the gamma vector and matroids	11
Lawrence polytopes and some invariants of a graph	11
Matroids and binary functions	12
Tutte polynomials and split matroids	12
Bounds on the number of cells of the Dressian	12
Filtrations of tope spaces of oriented matroids	13
Contracting a Single Element in a Transversal Matroid	13
Cyclic matroids	14
Frame matroids with a distinguished frame element	14
Piecing together quasi-graphic matroids from graphic pieces	14
Matroids of Symmetric Rigid Frameworks	15
A coarse Erdős-Pósa theorem	15
Erdős-Pósa property of tripods in directed graphs	16
Reconfiguration of Basis Pairs in Regular Matroids	16
JASMINE HALL	17
Problems on Group-labeled Matroid Bases	17

45

Geometric methods for matroid theory

Author: Graham Denham¹

¹ *Western University*

Corresponding Author: gdenham@uwo.ca

Convex geometry has played a role in matroid theory for at least fifty years, while toric and tropical geometry have been more recent contributors. Results over the last decade have highlighted the power of various geometric or geometrically inspired techniques in matroid theory. I will survey some recent progress in this direction, with the aim of strengthening connections with other aspects of matroid theory.

5

Is There Hope?

Author: Geoff Whittle¹

¹ *Victoria University*

Corresponding Author: geoffwhittle201@gmail.com

We know the excluded minors for the matroids representable over $\text{GF}(2)$, $\text{GF}(3)$ and $\text{GF}(4)$. What are the chances of explicitly finding the excluded minors for matroids representable over $\text{GF}(5)$?

We have an excellent understanding of the classes that arise when we consider matroids representable over $\text{GF}(2)$ and other fields. We also have an algebraic understanding of the classes that arise for matroids representable over $\text{GF}(3)$ and other fields. What are the chances of describing the classes for $\text{GF}(4)$ and other fields?

Recent explicit descriptions of the excluded minors for the class of 2-regular matroids and 3-regular matroids (almost) can be thought of as baby steps towards the resolution of the questions posed above. But let's be real, is there genuine hope?

34

Characterising matroids representable over all fields of size at least four

Author: Nick Brettell¹

¹ *Victoria University of Wellington*

Corresponding Author: nick.brettell@vuw.ac.nz

A matroid is regular if it is representable over every field. Characterisations of regular matroids are known (Tutte, 1958) in terms of excluded minors, having a totally unimodular representation, and representability over $\text{GF}(2)$ and some field of characteristic not two. A matroid is near-regular if it is representable over every field of size at least three. Characterisations of near-regular matroids are known in terms of excluded minors (Hall, Mayhew, van Zwam, 2011), having a near-unimodular representation, and representability over each of $\{\text{GF}(3), \text{GF}(8)\}$, or each of $\{\text{GF}(3), \text{GF}(4), \text{GF}(5)\}$ (Whittle 1997). In this talk, we consider matroids representable over all fields of size at least four. None of these three types of characterisation are known for this class, but I'll discuss some partial progress

for each, and some related open problems. This will include joint work with James Oxley, Charles Semple, and Geoff Whittle; Rudi Pendavingh; and Rutger Campbell.

25

On the matroid representation over finite rings

Author: Koji Imamura¹

Co-author: Keisuke Shiromoto¹

¹ *Kumamoto University*

Corresponding Author: imamura.math.engr@gmail.com

Matroids introduced by H. Whitney abstract some properties of linear independence among vectors in a vector space over a field. Nevertheless, it is well-known that almost all matroids are non-representable as a set of vectors over a field. It is one of the most significant problems to determine whether a given matroid is representable over some field. In this talk, we propose some representations of matroids by using matrices over finite rings. We will provide a method to construct a matroid by a matrix over a finite ring and some conditions for a matrix over a finite ring to yield some matroid. We also show that some well-known non-representable matroids can be represented by a matrix over a finite ring by our construction.

46

Towards Rota's conjecture for gain-graphic matroids

Author: Dillon Mayhew¹

Co-author: Daryl Funk²

¹ *University of Leeds*

² *Douglas College*

Corresponding Author: dillonmayhew@gmail.com

Rota's famous conjecture for representable matroids says that when F is a finite field, there are only finitely many minimal obstructions for the class of matroids that are linearly-representable using vectors with F as the field of scalars. A proof has been announced by Geelen, Gerards, and Whittle.

Gain-graphic matroids are analogues to matroids represented by vectors: instead of representing the matroid using numbers from a field, we use elements from a group. In order to represent a matroid using group elements, we require an intermediate object known as a gain-graph. Multiple theorems show us that finite-field-representable matroids and finite-gain-graphic matroids play symmetric roles in structural matroid theory: if we want to understand the structure of minor-closed families of matroids, we must understand both representable and gain-graphic classes. So it is natural to seek an analogue of Rota's conjecture for gain-graphic matroids: when H is a finite group, there are only finitely many minimal obstructions for the class of H -gain-graphic matroids.

In this talk I will outline our intended path towards Rota's conjecture for gain-graphic matroids. This is joint work with Daryl Funk.

3

Rigidity and reconstruction in matroids of highly connected graphs

Author: Dániel Garamvölgyi¹

¹ *Alfréd Rényi Institute of Mathematics, Budapest*

Corresponding Author: daniel.garamvolgyi@ttk.elte.hu

A *graph matroid family* \mathcal{M} is a family of matroids $\mathcal{M}(G)$ defined on the edge set of each finite graph G in a compatible and isomorphism-invariant way. We say that \mathcal{M} has the *Whitney property* if there is a constant c such that every c -connected graph G is uniquely determined by $\mathcal{M}(G)$. Similarly, \mathcal{M} has the *Lovász-Yemini property* if there is a constant c such that for every c -connected graph G , $\mathcal{M}(G)$ has maximal rank among graphs on the same number of vertices.

In the talk, I will discuss old and new results about these notions. In particular, we will see that for “most” graph matroid families, the Whitney and the Lovász-Yemini properties are equivalent. I will also talk about how related ideas led to solutions of conjectures of Kriesell and Thomassen on highly connected graphs. (Partly based on joint work with Tibor Jordán, Csaba Király and Soma Villányi.)

40

Infinite matroids on lattices

Author: Peter Nelson¹

Co-author: Andrew Fulcher²

¹ *University of Waterloo*

² *University College Dublin*

Corresponding Author: apnelson@uwaterloo.ca

There are at least two well-studied ways to extend matroids to more general objects - one can allow the ground set to be infinite, or instead define the concept of independence on a lattice other than a set lattice. I will give several cryptomorphic definitions of an object that generalizes a matroid in both these ways at once, and argue that they are (in some ways) nicer than the usual finite matroid axioms. This is joint work with Andrew Fulcher.

43

Unbounded matroids, polymatroids and subspace arrangements

Author: José Samper¹

¹ *Pontifical Catholic University of Chile*

Corresponding Author: jsamper@uc.cl

The goal of the talk is to introduce the theory of unbounded matroids. These are possibly unbounded Polyhedra in \mathbb{R}^d whose vertices are 01 whose edge directions are of the form $e_i - e_j$. We will explore some cryptomorphic variations of these definitions involving linear extensions of posets and relate them to matroids and polymatroids. We will discuss how these objects can be used to study the geometry of subspace arrangements and present several questions inspired by classical results for matroids. Based on joint work with J. Berggren and J. Martin.

7

Tropical Ideals

Author: Felipe Rincon^{None}

Corresponding Author: f.rincon@qmul.ac.uk

Tropical ideals are combinatorial objects introduced with the aim of giving tropical geometry a solid algebraic foundation. They can be thought of as combinatorial generalizations of the possible collections of subsets arising as the supports of all polynomials in an ideal of the polynomial ring. In general, the structure of tropical ideals is dictated by a sequence of ‘compatible’ matroids. In this talk I will introduce and motivate the notion of tropical ideals, explain their connection to matroid symmetric powers, and discuss work studying some of their main properties and their possible associated varieties.

11

Amalgams of matroids, fibre products and tropical graph correspondences

Author: Dmitry Mineev¹

¹ *Bar-Ilan University, Ramat Gan, Israel*

Corresponding Author: dmitry.mineyev@gmail.com

We discuss the interplay between certain properties of pairs of matroids and intersection theory of their Bergman fans, which has been extensively studied during the last decade. This provides another connection between tropical algebraic geometry and combinatorics of matroids.

In the first part of the talk, we formulate the theorem claiming that the proper amalgam of two matroids M_1 and M_2 along their common restriction N exists if and only if the tropical fibre product of Bergman fans $B(M_1) \times_{B(N)} B(M_2)$ is positive. In the second part of the talk, we introduce the notion of tropical correspondence. These correspondences are tropical subcycles in the product of Bergman fans, similar to their analogues in algebraic geometry. We define a “graph correspondence” for the maps of lattices of flats and prove that this construction is functorial in the case of “covering” maps of lattices. This is done via explicit combinatorial description involving generalization of Bergman fan which we name a “flag fan”.

We conclude with an overview of the open questions that arise from this perspective.

37

Speyer’s g conjecture via external activity of a pair of matroids

Author: Alex Fink¹

¹ *Queen Mary University of London*

Corresponding Author: a.fink@qmul.ac.uk

In 2009, looking to bound the face vectors of matroid subdivisions and tropical linear spaces, Speyer introduced the g -invariant of a matroid. He proved its coefficients nonnegative for matroids representable in characteristic zero and conjectured this in general. Later, Shaw and Speyer and I reduced the question to positivity of the top coefficient. In work in progress with Berget, we prove the conjecture. This talk will overview the combinatorial ingredients of the proof. The principal ones are the restriction to the diagonal D of the Dilworth truncation of a direct sum of two matroids, and a generalisation of external activity for D using the two matroids it is constructed from.

1

Symmetric tropical rank 2 matrices

Author: Kisun Lee¹

¹ *Clemson University*

Corresponding Author: kisunl@clemson.edu

In this talk, we discuss the tropicalization of the space of symmetric rank two matrices. Analogous to the result of Markwig and Yu for general tropical rank two matrices, it has a simplicial complex structure as the space of symmetric bicolored trees. As properties of this simplicial complex, we point out that this space is shellable and give several descriptions of the algebraic matroid of the space of symmetric rank two matrices. This is a joint work with May Cai and Josephine Yu.

23

Baker-Bowler theory for Lagrangian Grassmannians

Author: Donggyu Kim¹

¹ *KAIST*

Corresponding Author: donggyu@kaist.ac.kr

Baker and Bowler (2019) showed that the Grassmannian can be defined over a tract, a field-like structure generalizing both partial fields and hyperfields.

This notion unifies theories for matroids over partial fields, valuated matroids, and oriented matroids. We extend Baker-Bowler theory to the Lagrangian Grassmannian which is the set of maximal isotropic subspaces in a $2n$ -dimensional symplectic vector space.

By Boege et al. (2019), the Lagrangian Grassmannian is parameterized into the projective space of dimension $2^{n-2}(4 + \binom{n}{2}) - 1$ and its image is cut out by certain quadrics.

We simplify a list of quadrics so that these are apparently induced by the Laplace expansions only concerning principal and almost-principal minors of a symmetric matrix.

From the idea that the strong basis exchange axiom of matroids captures the combinatorial essence of the Grassmann-Plücker relations, we define matroid-like objects, called antisymmetric matroids, from the quadrics for the Lagrangian Grassmannian.

We also provide its cryptomorphic definition in terms of circuits capturing the orthogonality and maximality of vectors in a Lagrangian subspace.

We define antisymmetric matroids over tracts in two equivalent ways, which generalize both Baker-Bowler theory and the parameterization of the Lagrangian Grassmannian.

It provides a new perspective on the Lagrangian Grassmannian over hyperfields such as the tropical hyperfield and the sign hyperfield.

Our proof involves a homotopy theorem for graphs associated with antisymmetric matroids, generalizing Maurer's homotopy theorem for matroids.

We also prove that if a point in the projective space satisfies the 3-/4-term quadratic relations for the Lagrangian Grassmannian and its supports form the bases of an antisymmetric matroid, then it satisfies all quadratic relations, which is motivated by the earlier work of Tutte (1958) for matroids and the Grassmannian.

27

Direct sum of q -matroids and linear sets

Author: Gianira Alfarano¹

¹ *University College Dublin*

Corresponding Author: gianira.alfarano@ucd.ie

q -Matroids, the q -analogue of matroids, have been intensively investigated in recent years in coding theory due to their close connection with rank metric codes. Indeed, in 2018 it was shown by Jurrius and Pellikaan that a rank metric code induces a q -matroid that captures many of the code's invariants. In this talk we will deal with the direct sum of q -matroids, a concept recently introduced by Ceria and Jurrius, with a particular focus on the question of representability. We will show that representing the direct sum of t uniform q -matroids is equivalent to constructing special linear sets which are almost scattered with respect to the hyperplanes. In addition, we will give explicit constructions of such linear sets, implying as a byproduct that the direct sum of uniform q -matroids is always representable. This is a joint work with Relinde Jurrius, Alessandro Neri and Ferdinando Zullo.

41

Almost all q -matroids are not representable

Author: Sebastian Degen¹

¹ *Bielefeld University*

Corresponding Author: sdegen@math.uni-bielefeld.de

The q -analogue of a combinatorial object arises by replacing finite sets with finite dimensional vector spaces. In particular we can view q -matroids as q -analogues of matroids. One motivation to study q -matroids stems from coding theory, as the representable q -matroids arise from rank-metric codes. In the matroidal setting Peter Nelson proved in 2018 that asymptotically almost all matroids are non-representable, therefore one can ask if the same holds true in the q -analogue. In this talk we investigate this question and provide a positive answer to it. For this purpose we give a lower bound on the number of all fixed dimensional q -matroids, using the theory of constant dimension codes and an upper bound on the number of all representable q -matroids, using the concept of zero patterns.

24

On a q -analogue of perfect matroid designs

Author: Shinya Kawabuchi¹

Co-author: Keisuke Shiromoto¹

¹ *Kumamoto University*

Corresponding Author: 230d8554@st.kumamoto-u.ac.jp

A Steiner system is characterized with the subsets (called blocks) satisfying that all t -subsets are included in exactly one block. A *perfect matroid design* (PMD) is a matroid whose flats of the same rank have the same size. Recently, E. Byrne et. al. proposed a q -analogue of PMDs (namely q -PMDs) and constructed a non-trivial q -PMD from a q -analogue of a Steiner system. In this talk, we show some properties of q -PMDs and demonstrate that a certain class of q -PMDs induces a q -Steiner system.

33

The cyclic flats of L-polymatroids

Author: Andrew Fulcher¹

Co-author: Eimear Byrne ¹

¹ *University College Dublin*

Corresponding Author: andrew.fulcher@ucdconnect.ie

In recent years, q-polymatroids have drawn interest because of their connection with rank-metric codes. For a special class of q-polymatroids called q-matroids, the fundamental notion of a cyclic flat has been developed as a way to identify the key structural features of a q-matroid. In this talk, we will see a generalisation of the definition of a cyclic flat that can apply to q-polymatroids, as well as a further generalisation, L-polymatroids. The cyclic flats of an L-polymatroid is essentially a reduction of the data of an L-polymatroid such that the L-polymatroid can be retrieved from its cyclic flats. As such, in matroid theory, cyclic flats have been used to characterise numerous invariants.

26

Graphs embedded on surfaces and their delta-matroids

Author: Carolyn Chun¹

Co-authors: Iain Moffatt ²; Ralf Rueckriemen ²; Steven Noble ³

¹ *University of Maryland*

² *Royal Holloway University of London*

³ *Birkbeck, University of London*

Corresponding Author: cchun@umd.edu

By “cutting out” the edges and vertices of a graph cellularly embedded in a surface, we obtain a ribbon graph. A partial Petrial is obtained from a ribbon graph by selecting a subset of the ribbon edges adding a half-twist to each. Delta-matroids generalize ribbon graphs similarly to the way that matroids generalize graphs. The ribbon graph partial Petrial is the analogue of a more general delta-matroid operation. In this talk, we characterize the set of delta-matroids that is closed under this delta-matroid operation by a set of minimal obstructions.

44

Graphs in surfaces, their one-face subgraphs, and the critical group

Author: Iain Moffatt¹

Co-authors: Criel Merino ²; Steven Noble ³

¹ *Royal Holloway University of London*

² *National Autonomous University of Mexico*

³ *Birkbeck, University of London*

Corresponding Author: iain.moffatt@rhul.ac.uk

Critical groups are groups associated with graphs. They are well-established in combinatorics; closely related to the graph Laplacian and arising in several contexts such as chip firing and parking functions. The critical group of a graph is finite and Abelian, and its order is the number of spanning

trees in the graph, a fact equivalent to Kirchhoff's Matrix–Tree Theorem.

What happens if we want to define critical groups for graphs embedded in surfaces, rather than for graphs in the abstract?

In this talk I'll offer an answer to this question. I'll describe an analogue of the critical group for an embedded graph. We'll see how it relates to the classical critical groups, as well as to Chumtsov's partial-duals, Bouchet's delta-matroids, and a Matrix–quasi-Tree Theorem of Macris and Pule.

This is joint work with Criel Merino and Steven D. Noble.

36

Clique minors in dense matroids

Author: Fernanda Rivera Omaña¹

Co-authors: Peter Nelson¹; Sergey Norin²

¹ *University of Waterloo*

² *McGill University*

Corresponding Author: mriveraomana@uwaterloo.ca

Given ℓ and t positive integers, we will give a singly exponential bound on the number of points a $U_{2,\ell}$ -minor-free matroid with can have without containing $M(K_t)$ as a minor.

18

Chordal matroids arising from generalized parallel connections

Authors: Dylan Douthitt¹; James Oxley¹

¹ *Louisiana State University*

Corresponding Author: jddouthitt@live.com

In 1961, Dirac showed that chordal graphs are exactly the graphs that can be constructed from complete graphs by a sequence of clique-sums. By analogy with Dirac's result, we define the class of $GF(q)$ -chordal matroids as those matroids that can be constructed from projective geometries over $GF(q)$ by a sequence of generalized parallel connections across projective geometries over $GF(q)$. We characterize the class of such matroids in terms of forbidden induced minors for all powers of q and also in terms of forbidden flats whenever q is prime. This talk is based on joint work with James Oxley.

22

Connectivity systems, path-width, and well-quasi-ordering

Author: Jim Geelen¹

Co-author: Rutger Campbell²

¹ *University of Waterloo*

² *Discrete Mathematics Group*

Corresponding Author: jfgeelen@uwaterloo.ca

We prove that, for any integer k , there are only finitely many excluded minors for the class of matroids of path-width k . This result is obtained as a consequence of a well-quasi-ordering result on connectivity systems of bounded path-width. This is joint work with Rutger Campbell.

19

Average Hyperplane-Size in Complex-Representable Matroids

Author: Matthew Kroecker¹

¹ *University of Waterloo*

Corresponding Author: mekroeke@uwaterloo.ca

In 1941, Melchior proved that the average size of a line in a simple rank-3 real-representable matroid is less than three. A similar theorem for the complex-representable matroids was proved by Hirzebruch in 1983. In this talk, we discuss the problem of extending these results to flats of higher rank. We show that, in every simple rank-4 real-representable matroid which is not the direct sum of two lines, the average size of a plane is at most an absolute constant. We also present a generalization of this result to hyperplanes of arbitrary rank.

This talk is based on joint work with Rutger Campbell, Jim Geelen and Ben Lund.

39

Convex geometry of building sets

Author: Rick Danner¹

¹ *University of Vermont*

Corresponding Author: richard.danner@uvm.edu

Building sets were introduced in the study of wonderful compactifications of hyperplane arrangement complements and were later generalized to finite meet-semilattices. Convex geometries, the duals of antimatroids, offer a robust combinatorial abstraction of convexity. Supersolvable convex geometries and antimatroids appear in the study of poset closure operators, Coxeter groups, and matroid activities. We prove that the building sets on a finite meet-semilattice form a supersolvable convex geometry. As an application, we demonstrate that building sets and nested set complexes respect certain restrictions of finite meet-semilattices unifying and extending results of several authors.

4

Combinatorics of boundary algebras.

Author: Jonathan Boretsky¹

Co-author: Jonah Berggren

¹ *MPI Leipzig*

Corresponding Author: jboresky@math.harvard.edu

Boundary algebras are an important tool in the categorification, by Jensen–King–Su and by Pressland, of cluster structures on positroid varieties, defined by Scott and by Galashin–Lam. Each connected positroid has a corresponding boundary algebra. We give a combinatorial description of the set of algebras which arise as the boundary algebra of some positroid. Using this combinatorial structure, we give the first complete description of the minimal relations in the boundary algebra. This talk is based on joint work with Jonah Berggren.

16

Complexity of log-concave inequalities in matroids

Author: Swee Hong Chan¹

Co-author: Igor Pak²

¹ *Rutgers University*

² *University of California, Los Angeles*

Corresponding Author: sweehong.chan@rutgers.edu

A sequence of nonnegative real numbers a_1, a_2, \dots, a_n , is log-concave if $a_i^2 \geq a_{i-1}a_{i+1}$ for all i ranging from 2 to $n - 1$. Log-concavity naturally arises in various aspects of mathematics, each characterized by different underlying mechanisms. Examples range from inequalities that are readily provable, such as the binomial coefficients $a_i = \binom{n}{i}$, to intricate inequalities that have taken decades to resolve, such as the number of independent sets a_i in a matroid M with i elements (otherwise known as Mason’s conjecture). It is then natural to ask if it can be shown that the latter type of inequalities is intrinsically more challenging than the former. In this talk, we provide a rigorous framework to answer this type of questions, by employing a combination of combinatorics, complexity theory, and geometry. This is joint work with Igor Pak and is intended for a general audience.

31

Chow rings and augmented Chow rings of matroids are equivariant γ -positive

Author: Hsin-Chieh Liao¹

¹ *University of Miami*

Corresponding Author: hxl752@miami.edu

Chow rings and augmented Chow rings of matroids were considered and played important roles in the settlement of the Heron–Rota–Welsh conjecture and the Dowling–Wilson top-heavy conjecture. Their Hilbert series have been extensively studied since then. It was shown by Ferroni, Mathern, Steven, and Vecchi, and independently by Wang, that the Hilbert series of Chow rings of matroids are γ -positive using inductive argument from semismall decompositions. However, they do not have an interpretation for the coefficients in the γ -expansion. In the recent paper, Angarone, Nathanson, and Reiner conjectured that Chow ring of matroids are equivariant γ -positive under the action of groups of automorphisms of matroids. We prove this conjecture, showing that both Chow rings and augmented Chow rings of matroids are equivariant γ -positive. Moreover, we obtain explicit descriptions for the coefficients of the equivariant γ -expansions.

38

Representations on the cohomology of the moduli space of pointed rational curves, permutation representations and log concavity

Author: Donggun Lee¹

Co-authors: Jinwon Choi ; Young-Hoon Kiem

¹ *Institute for Basic Science*

Corresponding Author: dglee@ibs.re.kr

The Chow ring of the graphical matroid of the complete graph with $n-1$ vertices (with respect to the minimal building set) is isomorphic to that of the moduli space of n -pointed rational curves. Therefore, it is naturally a graded representation of the n -th symmetric group.

In this talk, we provide a closed-form formula for this representation, in terms of weighted rooted trees, which we introduce as combinatorial gadgets. Using this, we explore the positivity of this representation as a permutation representation, and the asymptotic log concavity of the multiplicities of the trivial representation. We will also discuss about the multiplicities of the other irreducible representations.

Based on joint works with Jinwon Choi and Young-Hoon Kiem.

2

Intrinsic properties of the gamma vector and matroids

Author: Soohyun Park¹

¹ *Hebrew University of Jerusalem*

Corresponding Author: soohyun.park@mail.huji.ac.il

We'll discuss the gamma vector, which is an invariant of (reciprocal) polynomials whose positivity lies somewhere between unimodality and real-rootedness. It shows up in many seemingly unrelated contexts including permutation statistics, signs of Euler characteristics vs. curvatures of (piecewise Euclidean) manifolds, and Chow rings of matroids. Before asking about positivity, a natural question to ask while trying to understand why this is the case is the question of *what* this invariant measures in the first place. While there are classes of objects collecting when positivity is observed (the second setting in particular), the invariant itself is rather poorly understood (except possibly the last component). We explore the question using an explicit formula for (an extension of) this invariant in terms of the input polynomial along with tools from algebraic geometry and geometric combinatorics. For example, we make observations that contrast with heuristics stated regarding the general lower bound behavior of h -vectors when it was originally defined. This perspective also leads to natural connections to convexity properties of polynomials specific to matroids and structures of matroid invariants coming from algebraic geometry.

30

Lawrence polytopes and some invariants of a graph

Author: Changxin Ding^{None}

We use two dual Lawrence polytopes P and P^* of a graph G to study the graph. The h -vector of the graphic (resp. cographic) matroid complex associated to G coincides with the h^* -vector of the

Lawrence polytope P (resp. P^*). In general, the h -vector is an invariant defined for an abstract simplicial complex, which encodes the number of faces of different dimensions. The h^* -vector, a.k.a. the δ -polynomial, is an invariant defined for a rational polytope obtained by dilating the polytope. By dissecting the Lawrence polytopes, we may study the h -vectors associated to the graph G at a finer level. In particular, we understand the reduced divisors of the graph G in a more geometric way.

32

Matroids and binary functions

Author: Graham Farr¹

¹ *Monash University*

Corresponding Author: graham.farr@monash.edu

A *binary function* is a function $f : 2^E \rightarrow \mathbb{C}$ for which $f(\emptyset) = 1$, where E is a finite ground set. Binary functions are closely related to quantum registers. They generalise binary matroids in the sense that any indicator function of a linear space over $\text{GF}(2)$ is a $\{0, 1\}$ -valued binary function (using the natural correspondence between subsets of E and their characteristic vectors in $\text{GF}(2)^E$); in fact, the author showed in 1993 that *every* matroid has an associated binary function, although it will not necessarily be just $\{0, 1\}$ -valued.

In a series of papers over 1993-2019, the author generalised some standard matroid transforms and operations —including rank, deletion, contraction, and duality (via the Hadamard transform) —to binary functions, and gave new versions of them that are parameterised by the complex numbers. In each of these settings, a theory of Tutte-Whitney polynomials was developed.

In this talk we review this work and discuss some recent results.

21

Tutte polynomials and split matroids

Author: Benjamin Schröter¹

¹ *KTH Royal Institute of Technology*

Corresponding Author: schrot@kth.se

Surprisingly many interesting invariants of graphs, hyperplane arrangements and matroids are valuate. The two most prominent examples of valuate matroid invariants are the Tutte polynomial and the \mathcal{G} -invariant. The relevance of the \mathcal{G} -invariant stems from its universal property that any other valuate invariant can be obtained as a specialization. Nevertheless, the most intensely studied invariant of matroids is clearly the Tutte polynomial as it respects deletion and contraction. An interesting question therefore is on which minor and duality closed classes of matroids is the Tutte polynomial universal. In my talk I will give answer to this question in which the well structured class of (elementary) split matroids plays a prominent role.

This talk is based on joint work with Luis Ferroni.

6

Bounds on the number of cells of the Dressian

Author: Rudi Pendavingh¹

¹ Eindhoven Technical University

Corresponding Author: r.a.pendavingh@tue.nl

A valuation of a matroid M with set of bases \mathcal{B} is a function $\nu : \mathcal{B} \rightarrow \mathbb{R}$ so that

- **(V)** for all $B, B' \in \mathcal{B}$ and all $e \in B \setminus B'$ there exists an $f \in B' \setminus B$ such that $\nu(B) + \nu(B') \geq \nu(B - e + f) + \nu(B' + e - f)$.

The Dressian $\mathcal{D}(M)$ is defined as the set of all valuations of M . The Dressian of M is the support of a polyhedral complex in $\mathbb{R}^{\mathcal{B}}$ whose cells are polyhedral cones. We investigate $\#\mathcal{D}(M)$, the number of cells, and $\dim \mathcal{D}(M)$, the maximum dimension of any cell.

We show that if M is a matroid on an n -element groundset of rank $r \geq t \geq 3$, then

$$\frac{\dim \mathcal{D}(M)}{\binom{n}{r}} \leq \frac{\max\{\dim \mathcal{D}(M/S), S \in \binom{E}{r-t} \text{ independent}\}}{\binom{n-r+t}{t}} \leq \frac{3}{n-r+3}$$

and

$$\frac{\log_2 \#\mathcal{D}(M)}{\binom{n}{r}} \leq \frac{\max\{\log_2 \#\mathcal{D}(M/S) : S \in \binom{E}{r-t} \text{ independent}\}}{\binom{n-r+t}{t}} O(\ln(n)) \leq \frac{O(\ln(n)^2)}{n}.$$

For uniform matroids $M = U(r, n)$ these upper bounds may be compared to the lower bounds

$$\frac{1}{n} \leq \frac{\dim \mathcal{D}(M)}{\binom{n}{r}}, \quad \frac{1}{n} \leq \frac{\log_2 \#\mathcal{D}(M)}{\binom{n}{r}}$$

that follow a previously known construction of valuations of $U(r, n)$ from matroids of rank r on n elements. For non-uniform matroids, we also obtain somewhat tighter upper bounds in terms of the number of free generators of the Tutte group of M .

Our methods include an analogue of Shearers' Entropy Lemma for bounding the dimension of a linear subspace, and a container method for collections of linear subspaces.

10

Filtrations of tope spaces of oriented matroids

Author: Chi Ho Yuen^{None}

Co-author: Kris Shaw¹

¹ University of Oslo

Corresponding Author: chyuen@math.nctu.edu.tw

Oriented matroids are matroids with extra sign data, and they are useful in the tropical study of real algebraic geometry. In order to study the topology of real algebraic/tropical hypersurfaces constructed from patchworking, Renaudineau and Shaw introduced an algebraically defined filtration of the tope spaces of oriented matroids (for a real hyperplane arrangement, the tope space is the homology of its complement) based on Quillen filtration. We will prove the equivalence of their filtration and the topologically defined Kalinin filtration, as well as the combinatorially defined Gelfand-Varchenko dual degree filtration.

15

Contracting a Single Element in a Transversal Matroid

Author: Sam Bastida¹

¹ *Victoria University of Wellington*

Corresponding Author: bastidsamu@myvuw.ac.nz

It is well known that the class of transversal matroids is not closed under contraction or duality. The complexity of deciding whether a minor or dual of a transversal matroid remains transversal is in Σ_2 and thus far there has been no improvement on this bound. We explore this issue, providing a polynomial time algorithm for determining whether a single element contraction of a transversal matroid remains transversal. If so, our algorithm also provides a transversal representation. We then develop the techniques used in search of a polynomial time algorithm for determining whether the dual of a transversal matroid remains transversal.

20

Cyclic matroids

Author: Charles Semple^{None}

Co-authors: Nick Brettell¹; Deborah Chun²; Tara Fife ; Gerry Toft

¹ *Victoria University of Wellington*

² *West Virginia University Institute of Technology*

Corresponding Author: charles.semple@canterbury.ac.nz

Tutte showed that wheels and whirls are precisely the 3-connected matroids in which every element is contained in a 3-element circuit and a 3-element cocircuit. In fact, wheels and whirls have the stronger property that there is a circular ordering of their ground set such that every two consecutive elements are contained in a 3-element circuit and a 3-element cocircuit. In this talk, we investigate matroids satisfying this property and discuss some recent results. This is joint work with Nick Brettell, Deborah Chun, Tara Fife, and Gerry Toft.

29

Frame matroids with a distinguished frame element

Author: James Davies¹

Co-authors: Cynthia Rodríguez ; Jim Geelen²

¹ *Cambridge*

² *University of Waterloo*

Corresponding Author: jgd37@cam.ac.uk

A matroid is frame if it may be extended such that it possesses a basis B (a frame) such that every element is spanned by at most two elements of B . This gives frame matroids natural graphical representations as biased graphs.

We characterise the inequivalent graphical representations of 3-connected frame matroids that have a fixed element ℓ in their frame B . One consequence is a polynomial time recognition algorithm for frame matroids with a distinguished frame element.

35

Piecing together quasi-graphic matroids from graphic pieces

Author: Nathan Bowler¹

Co-authors: Daryl Funk²; James Davies³; Jim Geelen⁴; Peter Nelson⁴

¹ *University of Hamburg*

² *Douglas College*

³ *Cambridge*

⁴ *University of Waterloo*

Corresponding Author: n.bowler1729@gmail.com

Quasi-graphic matroids provide a common generalisation of frame and lifted graphic matroids, with better algorithmic properties. We explain how to efficiently construct a framework for a quasi-graphic matroid given a schema for dividing that framework into pieces, each of which is graphic. This is joint work with James Davies, Daryl Funk, Jim Geelen and Peter Nelson.

14

Matroids of Symmetric Rigid Frameworks

Author: Katie Clinch¹

¹ *UNSW Sydney*

Corresponding Author: katharine.clinch@gmail.com

A 2-dimensional framework (G, p) is an embedding of a graph $G = (V, E)$ into 2-dimensional space, such that each vertex v is assigned a pair of coordinates $p(v) = (x_v, y_v) \in \mathbb{R}^2$. A framework is rigid if the only motions of the vertices in the plane that preserve all edge lengths are isometries of the plane. It is well-known (since at least the 1920s) that when the set of coordinates $p(V)$ are algebraically independent over \mathbb{Q} , then the rigidity of the framework is determined by the rank of the rigidity matroid of the graph.

However, real-life structures are typically not generic, and often exhibit symmetry. In the last 30 years there has been a lot of research into extending the definition of the rigidity matroid to different symmetry groups. Many of these results first model the framework as a group-labelled graph and then define a matroid on the graph. In this talk, I'll present an approach that unifies many of these results. Unfortunately the main open problem in the area, characterising rigidity under even dihedral groups, still remains open.

28

A coarse Erdős-Pósa theorem

Authors: Jungho Ahn¹; Jochen Pascal Gollin²; Tony Huynh³; O-joung Kwon⁴

¹ *KIAS*

² *Institute for Basic Science*

³ *Sapienza University of Rome*

⁴ *Hanyang university*

Corresponding Author: ojoungkwon@hanyang.ac.kr

An induced packing of cycles in a graph is a set of vertex-disjoint cycles with no edges between them. We generalise the classic Erdős-Pósa theorem to induced packings of cycles. More specifically, we show that there exists a function $f(k) = \mathcal{O}(k \log k)$ such that for every positive integer k , every graph G contains either an induced packing of k cycles or a set X of at most $f(k)$ vertices such that the closed neighbourhood of X intersects all cycles in G . Our proof is constructive and yields a polynomial-time algorithm finding either the induced packing of cycles or the set X . Furthermore, we show that for every positive integer d , if a graph G does not contain two cycles at distance more than d , then G contains sets $X_1, X_2 \subseteq V(G)$ with $|X_1| \leq 12(d+1)$ and $|X_2| \leq 12$ such that after removing the ball of radius $2d$ around X_1 or the ball of radius $3d$ around X_2 , the resulting graphs are forests.

As a corollary, we prove that every graph with no $K_{1,t}$ induced subgraph and no induced packing of k cycles has tree-independence number at most $\mathcal{O}(tk \log k)$, and one can construct a corresponding tree-decomposition in polynomial time. This resolves a special case of a conjecture of Dallard et al. (arXiv:2402.11222), and implies that on such graphs, many NP-hard problems, such as Maximum Weight Independent Set, Maximum Weight Induced Matching, Graph Homomorphism, and Minimum Weight Feedback Vertex Set, are solvable in polynomial time. On the other hand, we show that the class of all graphs with no $K_{1,3}$ induced subgraph and no two cycles at distance more than 2 has unbounded tree-independence number.

This is joint work with Jungho Ahn, Pascal Gollin, and Tony Huynh.

47

Erdős-Pósa property of tripods in directed graphs

Author: Meike Hatzel^{None}

Co-authors: Karolina Okrasa ; Marcin Briański ; Michał Pilipczuk

Corresponding Author: research@meikehatzel.com

Let D be a directed graphs with distinguished sets of sources $S \subseteq V(D)$ and sinks $T \subseteq V(D)$. A *tripod* in D is a subgraph consisting of the union of two S - T -paths that have distinct start-vertices and the same end-vertex, and are disjoint apart from sharing a suffix.

This talk presents a proof that tripods in directed graphs exhibit the Erdős-Pósa property. More precisely, there is a function $f: \mathbb{N} \rightarrow \mathbb{N}$ such that for every digraph D with sources S and sinks T , if D does not contain k vertex-disjoint tripods, then there is a set of at most $f(k)$ vertices that meets all the tripods in D .

One of the tools applied to obtain this result is the matroid intersection theorem for gammoids. The presented work is joint with Marcin Briański, Karolina Okrasa and Michał Pilipczuk.

17

Reconfiguration of Basis Pairs in Regular Matroids

Authors: Kristóf Bérczi¹; Bence Mátravölgyi²; Tamás Schwarcz¹

¹ ELTE Eötvös Loránd University

² ETH Zürich

Corresponding Author: tamas.schwarcz@ttk.elte.hu

In recent years, combinatorial reconfiguration problems have attracted great attention due to their connection to various topics such as optimization, counting, enumeration, or sampling. One of the most intriguing open questions concerns the exchange distance of two matroid basis sequences, a problem that appears in several areas of computer science and mathematics. White (1980) proposed a conjecture for the characterization of two basis sequences being reachable from each other by symmetric exchanges, which received a significant interest also in algebra due to its connection to toric ideals and Gröbner bases. In this work, we verify White's conjecture for basis sequences of length two in regular matroids, a problem that was formulated as a separate question by Farber, Richter, and Shank (1985) and Andres, Hochstättler, and Merkel (2014). Most of previous work on White's conjecture has not considered the question from an algorithmic perspective. We study the problem from an optimization point of view: our proof implies a polynomial algorithm for determining a sequence of symmetric exchanges that transforms a basis pair into another, thus providing the first polynomial upper bound on the exchange distance of basis pairs in regular matroids. As a byproduct, we verify a conjecture of Gabow (1976) on the serial symmetric exchange property of matroids for the regular case.

42

JASMINE HALL

Author: Jasmine Hall¹

¹ *Victoria University of Wellington*

Corresponding Author: jasmine.hall@vuw.ac.nz

TBD

13

Problems on Group-labeled Matroid Bases

Authors: Florian Hörsch¹; András Imolay²; Ryuhei Mizutani³; Taihei Oki³; Tamás Schwarcz²

¹ *CISPA*

² *ELTE Eötvös Loránd University*

³ *The University of Tokyo*

Corresponding Author: oki@mist.i.u-tokyo.ac.jp

Consider a matroid equipped with a labeling of its ground set to an abelian group. We define the label of a subset of the ground set as the sum of the labels of its elements. We study a collection of problems on finding bases and common bases of matroids with restrictions on their labels. For zero bases and zero common bases, the results are mostly negative. While finding a non-zero basis of a matroid is not difficult, it turns out that the complexity of finding a non-zero common basis depends on the group. Namely, we show that the problem is hard for a fixed group if it contains an element of order two, otherwise it is polynomially solvable.

As a generalization of both zero and non-zero constraints, we further study F -avoiding constraints where we seek a basis or common basis whose label is not in a given set F of forbidden labels. Using algebraic techniques, we give a randomized algorithm for finding an F -avoiding common basis of two matroids represented over the same field for finite groups given as operation tables. The study of F -avoiding bases with groups given as oracles leads to a conjecture stating that whenever an F -avoiding basis exists, an F -avoiding basis can be obtained from an arbitrary basis by exchanging at most $|F|$ elements. We prove the conjecture for the special cases when $|F| \leq 2$ or the group is ordered. By relying on structural observations on matroids representable over fixed, finite fields, we verify a relaxed version of the conjecture for these matroids. As a consequence, we

obtain a polynomial-time algorithm in these special cases for finding an F -avoiding basis when $|F|$ is fixed.