Contribution ID: 30

Type: not specified

## Freeze-in Inelastic Dirac Dark Matter and H0 tension

Tuesday, 23 July 2024 16:10 (25 minutes)

The discrepancy in the estimation of the Hubble constant, as measured by the local distance ladder and the Cosmic Microwave Background (CMB) observations, has become a significant concern, reaching a  $4\sigma$  to  $6\sigma$  level of significance. In this study, we investigate a particle physics model featuring an inelastic Dirac dark matter framework with a gauged  $U(1)_D$  symmetry, aimed at mitigating this tension in a cosmologically consistent manner while maintaining feeble coupling between the dark and visible sectors. Our model proposes the production of dark sector particles through a freeze-in mechanism, with the two sectors retaining distinct temperatures due to minimal kinetic mixing. Leveraging the presence of a light mediator, the dark matter particles undergo late-stage annihilation into these mediators via resonantly enhanced Sommerfeld cross-sections. Subsequently, these mediators annihilate into self-interacting dark radiations, selectively influencing the dynamics during the CMB era without perturbing Big Bang Nucleosynthesis (BBN), thus offering a dual solution by alleviating both the Hubble constant ( $H_0$ ) and  $\sigma_8$  tensions concurrently.

**Presenter:** Mr CHO, Wonsub (SKKU)

Session Classification: Presentations