




THE MORA PROJECT



# INPC 2025

May 25-30, 2025  
DCC, Daejeon, Korea



# Current status of MORA at IGISOL

Luis Miguel Motilla Martinez, on behalf of the MORA collaboration  
INPC2025



UNIVERSITÉ  
CAEN  
NORMANDIE






# Contents

- CP Violation and D correlation, the search for baryogenesis
- MORA Setup, IGISOL
- Latest experimental developments
- Outlook and conclusions


# CP Violation and D correlation, baryogenesis



# In search of CP Violation

- **Matter** and **antimatter** should have been created in **equal** amounts
- Why do we **only** see **baryonic matter**?
- Sakharov **conditions** for **baryogenesis**
  - Baryon number violation
  - **CPV**
  - Process out of equilibrium

- **CPV** known in the **SM**
  - Seen in decay of **K**, **B** and **D** mesons
  - **Not enough** to explain baryogenesis
- 
- **Where to find CPV**
    - **In the SM**: complex phase **CKM matrix**
    - **Outside SM**: D-correlation of beta decay



# $\beta$ decay D-Correlation

In a **GT-F** mixed  $\beta$ -decay the energy phase space can be written as \*

$$\omega(\langle J \rangle | E_e, \Omega_e, \Omega_\nu) dE_e d\Omega_e d\Omega_\nu = \frac{F(\pm Z, E_e)}{(2\pi)^5} p_e E_e (E^0 - E_e)^2 dE_e d\Omega_e d\Omega_\nu$$

$$\times \xi \left\{ 1 + a \frac{\mathbf{p}_e \cdot \mathbf{p}_\nu}{E_e E_\nu} + b \frac{m}{E_e} + c \left[ \frac{\mathbf{p}_e \cdot \mathbf{p}_\nu}{3E_e E_\nu} - \frac{(\mathbf{p}_e \cdot \mathbf{j})(\mathbf{p}_\nu \cdot \mathbf{j})}{E_e E_\nu} \right] \left[ \frac{J(J+1) - 3\langle (\mathbf{J} \cdot \mathbf{j})^2 \rangle}{J(2J-1)} \right] \right.$$

$$\left. \frac{\langle \mathbf{J} \rangle}{J} \cdot \left[ A \frac{\mathbf{p}_e}{E_e} + B \frac{\mathbf{p}_\nu}{E_\nu} + D \frac{\mathbf{p}_e \times \mathbf{p}_\nu}{E_e E_\nu} \right] \right\}$$

$\beta$  asymmetry


$\beta$ - $\nu$  correlation

Fierz interference

D correlation

**D** is **non-zero** for **T** reversal violation

\* Jackson, J. D.; Treiman, S. B.; Wyld, H. W. (1957) *Phy Rev* 106(3), 517-521



A precise measurement of **D** would allow to:

- Give **direct constraints** on CP-violating **Wilson coefficients** in the nucleon-level EFT
- **Probe specific BSM Models**, like **L-R symmetric** model and **Leptoquark** model
- Measure **Final State Interactions**, predicted by the SM

## Determination of the D-Correlation

As previously said: we need a **GT-F** mixed decay and **polarization of the ions**

→ We need to maximize  $F(X)$  and the polarization degree of ions to get the highest sensitivity for CPV

$$D \equiv \sin(\varphi_{AV}) \cdot \overbrace{\frac{2\rho}{1+\rho^2} \cdot \left(\frac{J}{J+1}\right)^{1/2}}^{F(X)}$$

→ Proportional to mixture degree and axial vector-vector phase

Final state interactions (**FSI**), never measured using D-correlation

From neutron measurement (emiT):  $\varphi_{AV} = 180.013^\circ \pm 0.028^\circ$  (68% CL)

*T. E. Chupp, R. L. Cooper, K. P. Coulter, et al. Phys. Rev. C 86, 035505 (2012)*

$$D_{FSI} \sim Z \alpha \frac{E_e}{M} \cdot A(\mu_f - \mu_i) \quad \text{Callan and Treiman, Phys. Rev. 162(1967)1494.}$$

# Selection of nuclei for D-Correlation

Maximal sensitivity to  $F(X)$

Maximal spin polarization

Two cases to separate new physics from FSI

Well produced  
in GANIL, easy  
to polarize,  
short half-life,  
GT-F decay

Shorter half-life,  
opposite sign  
for FSI, good  
candidate for  
IGISOL

	neutron	$^{19}\text{Ne}$	$^{23}\text{Mg}$	$^{35}\text{Ar}$	$^{39}\text{Ca}$
Sensitivity $F(X)$	0,43	-0,52	-0,65	0,41	0,71
$D_1 (\times 10^{-4})$	0,11	2,31	2,64	0,43	-0,47
$D_2 (\times 10^{-4})$	0,02	0,17	0,16	0,01	-0,02


$$D_{\text{neutron}} = (-0.94 \pm 1.89 \pm 0.97) \cdot 10^{-4} \quad D_{^{19}\text{Ne}} = (1 \pm 6) \cdot 10^{-4}$$

$$D_{\text{FSI}}(p_e) = \left( D_1 \cdot \frac{p_e}{p_{\text{emax}}} + D_2 \cdot \frac{p_{\text{emax}}}{p_e} \right) \times 10^{-4}$$

Best measurement so far, *statistics limited*, but not enough for CPV

Callan and Treiman, Phys. Rev. 162(1967)1494.  
Chen, Phys. Rev. 185(1969)2003.

# Laser polarization of $^{23}\text{Mg}$




- Pump the ions to  $M_F = \pm 2$  for spin J polarization
- We use a **dipole** excitation with a **circular polarized laser** to **increase(decrease)** the **projection number  $M_F$**
- $^{39}\text{Ca}$  has a more complex structure to polarize than  $^{23}\text{Mg}$  (2 lasers vs 1 laser)
- $^{23}\text{Mg}$  is the **best candidate for the first experiments at IGISOL**
  - First candidate for DESIR
  - Easy to produce and polarize
  - High sensitivity to FSI and new physics




---

# MORA Setup, IGISOL



# MORA Trap


Open Paul Trap developed in LPC Caen



Paul Trap consisting on 3 pairs of **electrodes** (R1-R6) and 2 **Einzel lenses**

The electrodes (R1-R6) trap the ions


Einzel lenses focus the beam



He gas is used to cool the ion cloud and optimize trapping efficiency


# MORA Setup

## Polarization measurement



- **Paul Trap** in the middle of the detection system
- 2 annular **Si detectors** in the line axis
  - For continuous polarization measurement by betas
- 4 **MCP detectors**, each 90° apart from each other
  - For recoil ion detection
- 4 **phoswich detectors**, between the MCP detectors
  - For beta detection
- Laser setup for polarization



## D correlation measurement



$$\frac{N_{coinc}^{+45^{\circ}} + N_{coinc}^{+135^{\circ}} - N_{coinc}^{-45^{\circ}} - N_{coinc}^{-135^{\circ}}}{N_{coinc}^{+45^{\circ}} + N_{coinc}^{+135^{\circ}} + N_{coinc}^{-45^{\circ}} + N_{coinc}^{-135^{\circ}}} = \delta \cdot D \cdot P$$



# Ion Production at IGISOL

- Radioactive ions produced by the **IGISOL** method. Collision of a **primary beam** (proton,  $^3\text{He}$ ...) with a **target** in a **gas cell**.
- Transfer reactions,  $^{24}\text{Mg}(p,d)$  or  $^{27}\text{Al}(^3\text{He},^4\text{He})$
- Ions transported within **purified He gas** into the **Sextupole Ion Guide (SPIG)**, later accelerated at **30keV** towards the **Dipole Magnet** mass separator





# Sextupole Ion Guide




RF sextupole ion guide, 3-5MHz ,  $\sim 1\text{kV } V_{pp}$

Transports ions from **high pressure** ( $\sim 0.1$  mbar)  
section into **high vacuum**



# IGISOL Beamline

1. Ion production
2. Isobar ( $A/q$ ) separation
3. Ions enter the HV cage, at 30kV
  - a. Ions get cooled and bunched in the RFQ Cooler Buncher
  - b. When exiting, they get accelerated to 2keV
4. Ions travel through the MRToF, where mass separation can be done
5. Arrival to MORA, at ground potential
  - a. Pulsed Drift Tube (PDT) to slow down and bring to ground potential the ions
  - b. Second PDT to further slow down the ions
  - c. MORA Trap and detection setup
  - d. The ions get ejected from the trap



---


# Latest developments

Laser polarization proof-of-principle

Beam purification efforts

# MORA Experiment at IGISOL

- Proton beam @30MeV hits the target,  $^{24}\text{Mg}(p,d)^{23}\text{Mg}$  reaction.
- Ions cool down and charge exchange to 1+, travel through SPIG and go to dipole magnet
- Choose Isobar ( $A/q=23$ ) and send beam to RFQ Cooler buncher
- Beam gets cooled and gets bunched for  $\sim 2.7\text{ms}$ . Sending one bunch to MORA every 11s.
- Bunched beam arrives to MORA PDT, where it gets pulsed down to -1kV to send to ground, ions at  $\sim 1\text{keV}$
- Bunched ions arrive to second PDT, where they get further deaccelerated to  $\sim 100\text{eV}$
- Ions get trapped for 6s
- After trap, the ions get ejected
- 5s of “off-trap” measurement for active background
- Measurement repeated for two different circular polarizations, changing every hour







# First conditions of MORA, Nov 2022


## Typical conditions of IGISOL

- He gas purified by zeolite cold trap and getter purified
- ~200mbar of He gas inside of gas cell
- Ti windows on the gas cell, nat Mg target facing the beam
- 5MHz SPIG RF, usual extraction voltages
- Usual Al ion guide

- Too high contamination, getting only ~10  $^{23}\text{Mg}$  ions per trap cycle, even with high production (>10k ions/s)
- Impossible to do polarization proof-of-principle
- The culprit:  $^{23}\text{Na}$  contamination
- The RFQ Cooler Buncher has a limited charge space, that gets saturated by too much  $^{23}\text{Na}^+$
- Since the problem is in the Cooler, we cannot make use of the MRToF for cleaning the beam
- Need to tackle contamination on the production

# MORA Timeline

Biggest challenge: Na contamination for trap efficiency





# In search of contamination's origin

## Possible origin: Target

- Traces of Na
- Nuclear reaction
- Surface ionization
- Sputtering

Various **target tests** done, **coatings** to prevent ionization and different **target material** (Al)

No improvement

## Possible origin: Gas cell

- Surface ionization / sputtering from the gas cell

**Changed** usual Al to graphite gas cell **baked** at **high temperature (1300°C)** for long period of time to **evaporate** traces

No improvement

## Possible origin: Gas

- Stopping He purification showed improvement
- Traces of Na coming from Zeolite
- Forming molecules with Na that could improve yield

Various **gas tests** done, different traces of gases over purified He and different **getter** and **cold trap** configurations

No improvement

# In search of contamination's origin

## Second half 2024



- **Progress made for polarization measurement**
  - Managed to get a **first measurement of polarization inversion** with  **$1.53\sigma$**  precision
- **Fixed PDT instabilities**
- Better understanding of the **source of contamination**
- **Only the SPIG can explain what we see**
  - Sputtering inside of the SPIG, affected by gas purity
  - Voltages greatly affect the contamination
  - Na only appears with beam on





## March 2025, Clean SPIG measurement



- Next step was **removal of Na from all SPIG surfaces**
- Used ACCLAB cooling loop water, conductivity of  $\sim 0.23 \mu\text{S}/\text{cm}$
- **SPIG disassembled, cleaned and reassembled** carefully to not spread contamination
- **Minimal ratio of Na:Mg yet**
- **Best results to this date!**
- The **rods** are the most **critical part**, cleaned with special care
- **New Nb rods** commissioned **baked at high temperature** for evaporating Na



# Polarization proof-of-principle

- Using detections ( $N_{\beta+}$ ) on the Si detectors (Si1 and Si2) in each polarization (+ or -,  $\uparrow$  or  $\downarrow$ ) to measure the asymmetry (A)
- $A = \alpha \cdot A_{\beta} \cdot P$ , where  $\alpha$  is a sensitivity factor estimated from simulations,  $A_{\beta}$  is the beta asymmetry and  $P$  the polarization degree
- Needed  $\sim 10^2$  ions per trap cycle to measure  $P$
- We need at least  $\sim 10^4$  ions per trap cycle to attempt to measure  $D$

$$A = \frac{N_{\beta+}^{\uparrow} - N_{\beta+}^{\downarrow}}{N_{\beta+}^{\uparrow} + N_{\beta+}^{\downarrow}} \propto A_{\beta} \cdot P$$



# March 2025, Results

Asymmetry Polarization  $\sigma^-$  :  $0.72 \pm 0.25$

Asymmetry Polarization  $\sigma^+$  :  $-0.42 \pm 0.16$

Asymmetry with full polarization of the cloud (from simulations):  $= 0.51 \pm 0.01$

$$A = 0.51 \pm 0.14$$


55% < P < 100% at 90% C.L.

Trapped ions/cycle

$90 \pm 9$  from Si detectors

$145 \pm 55$  from coincidences

$$A = \frac{N_{\beta+}^{\uparrow} - N_{\beta+}^{\downarrow}}{N_{\beta+}^{\uparrow} + N_{\beta+}^{\downarrow}} \propto A_{\beta} \cdot P$$



## Conclusions




Recent developments on MORA permitted to **fix old problems of trapping stability** and reduce the **beam contamination** issue, allowing us to have a **first measurement of the polarization** proof-of-principle

**Na contamination got low enough** that we can use the **MRTof** to **precisely measure the yield and contaminants**, and explore how to **improve** them.

In the **upcoming beamtime**, we will use the **new SPIG rods (Nb baked at high temperature)**, the contamination analysis and everything we have learned along the way, we will **measure the polarization degree  $P$  with higher precision**, and **start the measurement of the  $D$  correlation**



# Thank you for your attention!



E. Liénard  
M. Benali  
V. Bosquet  
S. Daumas-Tschopp  
L. Hayen  
Y. Merrer  
X. Fléchar  
G. Quémener  
A. De Roubin



M. Gonzalez-Alonso



A. Falkowski  
A. Rodriguez – Sanchez



M.L. Bissel



**P. Delahaye - P.I.**  
F. De Oliveira  
S. K. Chinthakayala  
C. Fougères  
G. Frémont  
N. Goyal  
N. Lecesne  
**M. JBayli**  
R. Leroy  
**L. M. Motilla**  
B.M. Retaillieu  
A. Singh  
J. C. Thomas



N. Severijns  
R.P. De Groote  
G. Neyens



I. Moore  
T. Eronen  
M. Reponen  
W. Gins  
V. Virtanen  
J. Romero  
A. Raggio  
A. Jaries  
A. Jokinen  
A. Kankainen  
S. Kujanpää  
M. Stryczyk  
S. Rinta-Antila  
Z. Ge




M. Kowalska






## Extra slides

# Sextupole Ion Guide



RF sextupole ion guide, 3-5MHz,  $\sim 1\text{kV } V_{pp}$

Transports ions from **high pressure** ( $\sim 0.1$  mbar) section into **high vacuum**



Source: ~~target?~~ gas cell? gas?

## Data Acquisition: June 2024

Installation of **baked C** gas cell trying to evaporate all Na → No immediate improvement.

**Second gas injection attempt, trying to find the gas that makes Na disappear**

Gases planned to be used: CO<sub>2</sub>, N<sub>2</sub>, He+CF<sub>4</sub>, SF<sub>6</sub>, He+Kr, He+Ne

Gases used: CO<sub>2</sub>, He+CF<sub>4</sub>, He+Kr

**Mass flow controller** allows for precise measurement of gas flow we are introducing into the chamber



Source: ~~target?~~ ~~gas cell?~~ gas?

## Data Acquisition: July 2024

- First polarization evidence ( $1.53\sigma$ ) measured
- The improvement was achieved mainly by having a **full dewar of LN** and **fine tuning the SPIG** voltages
- If the Na was produced at a lower energy/other mechanism, it is safe to assume that the extraction would be much different
- Further **gas injection test** did not prove **useful**, but it might be coming from the zeolite in the purification system
- PDT stability issues and how to fix them



Ringin from the PDT after pulsing down







Source: ~~target?~~ ~~gas?~~ cell? gas?

## Data Acquisition: Dec2024

- We used charcoal and SiO<sub>2</sub> based cold trap to avoid any zeolite, and new filters inside of LN to simulate the events in July (full dewar)
- Immediately we saw **no improvement** in the yield Na:Mg
- **The only suspect left was the SPIG**
- The SPIG is coherent with all other observations:
  - Effects of gas purity/pressure
  - The Na<sup>+</sup> can be trapped in the SPIG
  - Na<sup>+</sup> only appears with beam on
- Fixed PDT!



# In search of contamination's origin

## July 2024 beamtime

- Observed **improvement** on ratio Na:Mg by completely **filling the cold trap**, getting the **filter submerged in LN**
- Instead of maximizing  $^{23}\text{Mg}$  yield, we tried to **minimize the contamination from  $^{23}\text{Na}$**  with the SPIG
- **Trace gas injection did not provide useful** for the Mg beam
- With this, we managed to get a **first measurement of polarization inversion** with  **$1.53\sigma$**  precision

## December 2024 beamtime

- Tried to replicate July condition, by using new cold traps with **activated charcoal** and  $\text{SiO}_2$
- **No improvement on contamination**, but the **PDT** was finally **fixed**
- After discarding other suspects, the only possibility left was the **SPIG as a source of Na**.




# Polarization proof-of-principle

- Using detections ( $N_{\beta+}$ ) on the Si detectors (Si1 and Si2) in each polarization (+ or -,  $\uparrow$  or  $\downarrow$ ) to measure the asymmetry (A)
- $A = \alpha \cdot A_{\beta} \cdot P$ , where  $\alpha$  is a sensitivity factor estimated from simulations,  $A_{\beta}$  is the beta asymmetry and  $P$  the polarization degree
- Needed  $\sim 10^2$  ions per trap cycle to measure  $P$
- We need at least  $\sim 10^4$  ions per trap cycle to attempt to measure  $D$

$$A = \frac{N_{\beta+}^{\uparrow} - N_{\beta+}^{\downarrow}}{N_{\beta+}^{\uparrow} + N_{\beta+}^{\downarrow}} \propto A_{\beta} \cdot P$$

- Using coincidences between Si and MCP to improve signal to noise ratio
- Then we need to use  $B_{\nu}$ , the neutrino asymmetry





## Data Analysis: Calibration

The calibration fit is done with a PENELOPE simulation

$\text{Bi}^{207}$  source for the Si detectors and 3 alpha sources for the RIDE detectors

Si calibration using 3 parameters and resolution

