



Contribution ID: 178

Type: **Contributed Oral Presentation**

Highly-Charged Radioactive Molecules: Amplifying Sensitivity for New Physics

Monday, 26 May 2025 14:15 (15 minutes)

One necessary extension to the Standard Model of Particle Physics (SM) [1-4] is one which describes the behaviour of the early universe that leads to the matter-antimatter asymmetry [5] which we observe today. It is commonly assumed that any explanation of this matter-antimatter imbalance must rely on the violation of the combined symmetry of charge conjugation (C) and parity (P) [6] that is presently, however, considered to be too weak in the SM. Thus, identifying a new source of CP violation is of critical importance. CP-violating effects, particularly those that originate within the atomic nucleus, can be investigated by combining precision techniques from atomic, molecular, and optical physics with rare isotopes produced at accelerator facilities such as TRIUMF in Canada [7]. In searches for CP-violating nuclear Schiff moments, for example, molecular systems offer a sensitivity advantage of 3-4 orders of magnitude [8-10] which can be further increased by up to a factor of 1000 when a radioactive, octopole-deformed nucleus is incorporated into the molecule [11]. Among these radionuclides, the short-lived protactinium isotope ^{229}Pa is thought to exhibit the highest sensitivity, once a suitable molecule has been identified. Recently, triply charged protactinium-monofluoride PaF^{3+} has been proposed as a highly attractive probe [12], which is isoelectronic to the well studied case of neutral RaF [13]. Experimentally, forming such 'highly charged' molecules remains a significant challenge, also because all other Pa isotopes are radioactive, complicating the development of the necessary techniques. In this talk, we present the experimental formation of stable, doubly-charged cerium monofluoride CeF^{2+} which is identified as an intriguing surrogate to PaF^{3+} with a very similar molecular structure. Specific sensitivities of CeF^{2+} to new physics also position the system as an interesting probe for disentangling sources of symmetry-violating behaviour inside the nucleus.

References

- [1] S. L. Glashow. Partial-symmetries of weak interactions. *Nuclear Physics*, 22:579–588, 1961.
- [2] S. Weinberg. A model of leptons. *Physical Review Letters*, 19:1264, 1967.
- [3] A. Salam. Weak and electromagnetic interactions. *Proceedings of the Eighth Nobel Symposium*, pages 367–377, 1968.
- [4] M. Veltman. Regularization and renormalization of gauge fields. *Nuclear Physics B*, 44:189–213, 1972.
- [5] L. Canetti, M. Drewes, and M. Shaposhnikov. Matter and antimatter in the universe. *New Journal of Physics*, 14:095012, 2012.
- [6] A.D. Sakharov. The initial stage of an expanding universe and the appearance of a nonuniform distribution of matter. *Soviet Physics JETP*, 22:241, 1966.
- [7] J. Dilling, R. Krücken, and G. Ball. Isac overview. *Hyperfine Interactions*, 225:1–8, 2014.
- [8] N.J. Fitch, J. Lim, E.A. Hinds, B.E. Sauer, and M.R. Tarbutt. Methods for measuring the electron's electric dipole moment using ultracold ybf molecules. *Quantum Science and Technology*, 6(1):014006, 2020.
- [9] ACME Collaboration, V. Andreev, D.G. Ang, D. DeMille, J.M. Doyle, G. Gabrielse, J. Haefner, N.R. Hutzler, Z. Lasner, C. Meisenhelder, B.R. O'Leary, C.D. Panda, A.D. West, E.P. West, and X. Wu. Improved limit on the electric dipole moment of the electron. *Nature*, 562:355–360, 2018.
- [10] T.S. Roussy, L. Caldwell, T. Wright, W.B. Cairncross, Y. Shagam, K.B. Ng, N. Schlossberger, S.Y. Park, A. Wang, J. Ye, and E.A. Cornell. An improved bound on the electron's electric dipole moment. *Science*, 381(6653):46–50, 2023.
- [11] R.G. Garcia Ruiz. Short-lived radioactive molecules: A sensitive laboratory for the study of fundamental symmetries. *APS Division of Nuclear Physics Meeting Abstracts*, 2019:KA-003, 2019.

[12] C. Zülch, R.F. Garcia Ruiz, S.M. Giesen, R. Berger, and K. Gaul. Cool molecular highly charged ions for precision tests of fundamental physics. Arxiv, arXiv: 2203.10333, 2022.

[13] S.M. Udrescu, S.G. Wilkins, A.A Breier, et al. Precision spectroscopy and laser-cooling scheme of a radium-containing molecule. Nature Physics, 20:202–207, 2024.

Primary author: SIMPSON, Rane (TRIUMF)

Co-authors: Mrs KWIATKOWSKI, Ania (TRIUMF); Mr ZUELCH, Carsten (University of Marburg); Mr CHARLES, Chris (TRIUMF); Ms BELOSEVIC, Ivana (ISOLDE); Mr NG, Kia Boon (TRIUMF); COLLABORATION, RadMol (TRIUMF); Mr BERGER, Robert (University of Marburg); Mr MALBRUNOT-ETTENAUER, Stephan (TRIUMF); COLLABORATION, TITAN (TRIUMF)

Presenter: SIMPSON, Rane (TRIUMF)

Session Classification: Parallel Session

Track Classification: Fundamental Symmetries and Interactions in Nuclei