





JAGIELLONIAN UNIVERSITY
IN KRAKOW



Search for the **Axion-Like-Particles** in the η meson decay with the **HADES** Detector

Marcin Zieliński

*M. Smoluchowski Institute of Physics
Jagiellonian University, Kraków, Poland*

 **INPC 2025**

Daejeon, Korea, May 26th, 2025



NATIONAL SCIENCE CENTRE
POLAND


General motivation

The general and main motivation for research is to answer the question:

Why does the Universe exist?

More specific question for physicists:

How did our 'Material Universe' survive the cooling after the Big Bang?



Big Bang:

an equal amount of matter and antimatter was produced during the hot phase

During cooling and expansion

matter and antimatter annihilated ☹️



Most of the cosmic energy/matter budget is of an unknown form


General motivation

Status of Standard Model in HEP:

- **The Standard Model has served us well for 50 years!**
- Recent measurements indicates SM can't be the final answer.
- Six categories of problems have arisen in SM.


Standard Model of Elementary Particles

three generations of matter (fermions)			interactions / force carriers (bosons)	
I	II	III		
mass $\approx 2.16 \text{ MeV}/c^2$ charge $\frac{2}{3}$ spin $\frac{1}{2}$ u up	mass $\approx 1.273 \text{ GeV}/c^2$ charge $\frac{2}{3}$ spin $\frac{1}{2}$ c charm	mass $\approx 172.57 \text{ GeV}/c^2$ charge $\frac{2}{3}$ spin $\frac{1}{2}$ t top	mass $\approx 125.2 \text{ GeV}/c^2$ charge 0 spin 0 H higgs	
mass $\approx 4.7 \text{ MeV}/c^2$ charge $-\frac{1}{3}$ spin $\frac{1}{2}$ d down	mass $\approx 93.5 \text{ MeV}/c^2$ charge $-\frac{1}{3}$ spin $\frac{1}{2}$ s strange	mass $\approx 4.183 \text{ GeV}/c^2$ charge $-\frac{1}{3}$ spin $\frac{1}{2}$ b bottom	mass 0 charge 0 spin 1 g gluon	
mass $\approx 0.511 \text{ MeV}/c^2$ charge -1 spin $\frac{1}{2}$ e electron	mass $\approx 105.66 \text{ MeV}/c^2$ charge -1 spin $\frac{1}{2}$ μ muon	mass $\approx 1.77693 \text{ GeV}/c^2$ charge -1 spin $\frac{1}{2}$ τ tau	mass 0 charge 0 spin 1 γ photon	
mass $< 0.8 \text{ eV}/c^2$ charge 0 spin $\frac{1}{2}$ ν_e electron neutrino	mass $< 0.17 \text{ MeV}/c^2$ charge 0 spin $\frac{1}{2}$ ν_μ muon neutrino	mass $< 18.2 \text{ MeV}/c^2$ charge 0 spin $\frac{1}{2}$ ν_τ tau neutrino	mass $\approx 91.188 \text{ GeV}/c^2$ charge 0 spin 1 Z Z boson	
			mass $\approx 80.3692 \text{ GeV}/c^2$ charge ± 1 spin 1 W W boson	



General motivation

Dark Sector Candidates, Anomalies, and Search Techniques



- In SM: violation from weak interaction is not sufficient to create observed asymmetry

- Parameter space for BSM is running out at HEP

- several anomalies in experiments point to possible new physics, weakly coupled to familiar matter in the 1 MeV - 1 GeV scale

Ref: Marco Battaglieri, [arXiv:1707.04591](https://arxiv.org/abs/1707.04591) [hep-ph]

Newest theoretical models prefer gauge bosons in MeV-GeV mass range as “...many of the more severe astrophysical and cosmological constraints that apply to lighter states are weakened or eliminated, while those from high energy colliders are often inapplicable” (B. Batell, M. Pospelov, A. Ritz – 2009)

How we can try to address SM problems experimentally?

- Searching for new particles
- Studying violation of discrete symmetries

**Axions and
Axion-Like-Particles**


Connection between Standard and Dark Matter

New Physics connects to Standard Model particles through four portals:

Portal	Particles	Operator(s)
“Vector”	Dark photons	$-\frac{\epsilon}{2\cos\theta_W} B_{\mu\nu} F'^{\mu\nu}$
“Axion”	Pseudoscalars	$\frac{a}{f_a} F_{\mu\nu} \tilde{F}^{\mu\nu}, \frac{a}{f_a} G_{i\mu\nu} \tilde{G}_i^{\mu\nu}, \frac{\partial_\mu a}{f_a} \bar{\psi} \gamma^\mu \gamma^5 \psi$
“Higgs”	Dark scalars	$(\mu S + \lambda S^2) H^\dagger H$
“Neutrino”	Sterile neutrinos	$y_N L H N$


Invisible, non-SM

Dark Matter production
Producing stable particles that could be (all or part of) Dark Matter




Visible, SM

Production of portal-mediators that decay to SM particles
Systematically exploring the portal coupling to SM particles



Mixed visible-invisible

Production of “rich” dark sectors
Testing the structure of the dark sector



Light mesons studies:

η/η'

Connection between Standard and Dark Matter

“Light dark matter must be neutral under SM charges, otherwise it would have been discovered at previous colliders”

[G. Krnjaic RF6 Meeting, 8/2020]

- The only known particles with all-zero quantum numbers: $Q = I = J = S = B = L = 0$ are the η/η' mesons and the Higgs boson (also the vacuum!) -> **very rare**
- The η meson is a Goldstone boson (the η' meson is not!)
- The η/η' decays are flavor-conserving reactions

Experimental advantages:

- Hadronic production cross section is quite large (~ 0.1 barn) \rightarrow much easier to produce than heavier mesons
 - All its possible strong decays are forbidden in lowest order by P and CP invariance, G-parity conservation and isospin and charge symmetry invariance.
 - EM decays are forbidden in lowest order by C invariance and angular momentum conservation
- Branching Ratio of processes from New Physics are enhanced compared to SM.

Axion-Like-Particles

How to explain observed anomalies
in view of existing experimental constraints for QCD Axion ?

Piophobic QCD axion

Must be short lived ($\sim 10^{-13}$ s) and decay predominantly to $e^+ e^-$

QCD Axion couples predominantly to the first generation of SM fermions (PQ charges vanish for second and third SM fermions)

The $a - \pi^0$ mixing at the level of $O(10^{-4})$

Then in SM Lagrangian axionic phases are directly ascribed to quark masses

$$m_u \rightarrow m_u e^{i\gamma^5 q_{PQ}^u a/f_a},$$

$$m_d \rightarrow m_d e^{i\gamma^5 q_{PQ}^d a/f_a},$$

$$m_e \rightarrow m_e e^{i\gamma^5 q_{PQ}^e a/f_a},$$

$$\frac{m_u}{m_d} \simeq \frac{Q_d^{PQ}}{Q_u^{PQ}} = \frac{1}{2}$$

$m_{u,d} \ll m_s$

isovector $\theta_{a\pi} \gg \theta_{a\eta}, \theta_{a\eta'}$

isoscalar $\theta_{a\pi} \ll \theta_{a\eta}, \theta_{a\eta'}$

Suppressed mixing-angle results in the isoscalar couplings of the axion

Axion-Like-Particles

Hadronic decay channels of η and η'
could be coupled to ALP's:

$$\eta \rightarrow \pi^+ \pi^- a \quad (\rightarrow e^+ e^-)$$


"axio-hadronic decay"

Using Resonance Chiral Theory (R χ T),
the low-lying resonances should be
included as degrees of freedom in the
R χ T Lagrangian




χ PT predictions for decay rates
significantly modified by inclusion
of resonance exchange.

*".... $O(10^{-2})$, is probably excluded or in tension
with observations but $O(10^{-4} - 10^{-3})$ likely
remains experimentally allowed, and within
the sensitivity."*



$$\text{BR}(\eta \rightarrow \pi\pi a) \sim 3 \cdot 10^{-8} - 4 \cdot 10^{-6}$$


Effects of pion-pion final-state interactions (FSI)



Axion-Like-Particles

Why previous measurements $\eta(\eta') \rightarrow \pi^+ \pi^- e^+ e^-$ did not see Axion signatures ?

Experiments up to now focused on studies of **CP invariance**:




Year	Exp.	Events number	Asymmetry	BR ($\eta \rightarrow \pi^+ \pi^- e^+ e^-$)
2009	KLOE-2	1555 ± 52	$(-0.6 \pm 2.5_{\text{stat}} \pm 1.8_{\text{syst}}) \times 10^{-2}$	$(2.68 \pm 0.09_{\text{stat}} \pm 0.07_{\text{syst}}) \times 10^{-4}$
2016	WASA-at-COSY	251 ± 17	$(-1.1 \pm 6.6_{\text{stat}} \pm 0.2_{\text{syst}}) \times 10^{-2}$	$(2.7 \pm 0.2_{\text{stat}} \pm 0.2_{\text{syst}}) \times 10^{-4}$
2007	WASA-CELSIUS	$16.3 \pm 4.9 \pm 2.0$	-----	$(4.3 \pm 1.3_{\text{stat}} \pm 0.4_{\text{syst}}) \times 10^{-4}$

Rejected events
 $m(e^+e^-) < 20 \text{ MeV}$

BES III – Result for the $\eta \rightarrow \pi^+ \pi^- e^+ e^-$ decay (also **CP invariance** studies):

R as a mas of axion



$$R^{\text{UP}} = \frac{\mathcal{B}(\eta \rightarrow \pi^+ \pi^- a) \cdot \mathcal{B}(a \rightarrow e^+ e^-)}{\mathcal{B}(\eta \rightarrow \pi^+ \pi^- e^+ e^-)}$$




$$(3.0 - 53.0) \times 10^{-3}$$


Ref: BESIII Col. arXiv:2501.10130v1 (2025)



JAGIELLONIAN UNIVERSITY
IN KRAKOW




HADES - High Acceptance Di-Electron Spectrometer





SIS 18

U ⁷³⁺	1.0 GeV/u	10 ⁹ ions/s
Protons	4.5 GeV	2.8x10 ¹³ /s
Pions	0.5-2 GeV/c	



➤ hades.gsi.de

HADES - High Acceptance Di-Electron Spectrometer





February 2022 measurement:

- proton – proton collisions at energy of $T = 4.5 \text{ GeV}$ using LH_2 target
- 28 days of measurement
- estimated total integrated luminosity
 $\sim 6.1 [\text{pb}^{-1}]$

- **START** – T0 reaction for ToF
- **RICH** – Cherenkov detector (di-electron e^+e^-)
- **MDC and STS** – track reconstruction
- **Magnet Coil** – generates magnetic field
- **ToF & RPC** – Time-of-Flight META detectors
- **ECAL** – electromagnetic calorimeter (photons)
- **Trigger logic based on InnerToF and Meta**

Lepton identification using HADES RICH Detector




$$m_{\pi} > m_e \quad \beta_{\pi} < \beta_e$$



$$\beta_{\pi} < \beta_{\text{Cher}} < \beta_e$$

$e^+ e^-$ creates
Cherenkov radiation

- Lepton identification base on signals in RICH.
- Threshold momentum for electrons 9 MeV and for pions 2500 MeV.
- Acceptance as a function of transverse momentum and $e^+ e^-$ invariant mass.



Particle selection and identification






Following particles have to be selected: $\pi^+ \pi^- e^+ e^-$

- leptons selected by correlation windows ($\theta_{\text{RICH}} - \theta_{\text{MDC}}$) in RICH and MDC
- pions selected by cuts on beta vs momentum distribution
- additional cuts for leptons: $\beta > 0.9$



Event selection for the $\eta \rightarrow \pi^+ \pi^- e^+ e^-$ decay

- vertexReco $z \in (-200 \text{ mm}, 0)$
- $\pi^+ \pi^- e^+ e^-$ missing mass vs inv. mass
(graphical cut)




Event selection for the $\eta \rightarrow \pi^+ \pi^- e^+ e^-$ decay

- $(e^+e^-)(\pi^+\pi^-)$ opening angle $< 50^\circ$ cut
- $\pi^+\pi^-$ invariant mass < 480 MeV cut




- $(e^+e^-)(\pi^+\pi^-)$ opening angle in CM $> 140^\circ$

$(e^+e^-)(\pi^+\pi^-)$ opening angle in C.o.M.




$(e^+e^-)(\pi^+\pi^-)$ opening angle in C.o.M.





Event selection for the $\eta \rightarrow \pi^+ \pi^- e^+ e^-$ decay

- all cuts were compared using $e^+e^-\pi^+\pi^-$ invariant mass
- Most of the multipion background was subtracted
- reduction of 86.78% events in total range of $e^+e^-\pi^+\pi^-$ invariant mass distribution (data)
- reduction of 10.16% events in η signal range (simulations)

The experimental invariant mass after the application of consecutive cuts.



Extraction of $\eta \rightarrow \pi^+ \pi^- e^+ e^-$ signal



Estimated number of signal events

2758

η peak mean (MeV)


548.40

η peak sigma (MeV)

32.59

Results

- Final distribution of e^+e^- invariant mass after background subtraction
- Estimated total efficiency and acceptance factor: $1.1 \cdot 10^{-3}$



Upper limit for the number of event estimated based on W. A. Rolke method:

Ref. *Nuc. Instr. and Met. in Phys. A*, 551, 2-3 (2012)

$$N_{\chi_{17}}^{\text{UL}} = 255 \text{ (CL=90\%)}$$

$$\text{BR}_{\eta \rightarrow \pi^+\pi^-\chi_{17}} < 2.58 \cdot 10^{-5}$$

$$\text{BR}^{\text{theory}}_{\eta \rightarrow \pi^+\pi^-\pi^0} < 4 \cdot 10^{-6}$$

Conclusions

- *η/η' mesons are an interesting place to look for dark particles because probe coupling to light quarks and gluons.*
- *Preliminary estimation of upper limit for the Axion-Like-Particle in decay $\eta \rightarrow \pi^+ \pi^- e^+ e^-$ is $BR < 2.58 \cdot 10^{-5}$.*

Further steps:

- Studies of systematical effects.
- More detailed simulations of η decays and background using transport models SMASH/GIBUU.
- Application of Machine Learning techniques (MVA, BDT) to reduce background.



Thank you for your attention!





Backup

Event selection for the $\eta \rightarrow \pi^+ \pi^- e^+ e^-$ decay

- Combinatorial background subtraction:

$$\langle N_{CB} \rangle = 2\sqrt{\langle N_{\pi^+\pi^-e^+e^-} \rangle \langle N_{\pi^+\pi^-e^-e^-} \rangle}$$




$$\langle N_{signal} \rangle = \langle N_{\pi^+\pi^-e^+e^-} \rangle - \langle N_{CB} \rangle$$




Simulations of signal and background

Signal and main background reactions:

- $pp \eta \rightarrow pp \pi^+ \pi^- e^+ e^-$
- $pp \eta \rightarrow pp \pi^+ \pi^- X17 (e^+ e^-)$
- $pp \eta \rightarrow pp \pi^+ \pi^- \pi^0 (e^+ e^- \gamma)$
- $pp \eta \rightarrow pp \pi^+ \pi^- \pi^0 (\gamma \gamma)$
- $pp \eta \rightarrow pp \pi^+ \pi^- \gamma$



Estimation of X17 contribution to signal region



Right peak (R)

$$\eta \rightarrow \pi^+ \pi^- e^+ e^-$$

$$\eta \rightarrow \pi^+ \pi^- \gamma$$

$$\eta \rightarrow \pi^+ \pi^- X17 (e^+ e^-)$$

Left peak (L)

$$\eta \rightarrow \pi^+ \pi^- \pi^0 (e^+ e^- \gamma)$$

$$\eta \rightarrow \pi^+ \pi^- \pi^0 (\gamma \gamma)$$

Reaction	Contribution	Branching ratio (BR)
$\eta \rightarrow \pi^+ \pi^- e^+ e^-$	39.95%	$2.68 \cdot 10^{-4}$
$\eta \rightarrow \pi^+ \pi^- X17 (e^+ e^-)$	26.28%	$1 \cdot 10^{-4}$
$\eta \rightarrow \pi^+ \pi^- \gamma$	33.77%	$4.28 \cdot 10^{-2}$


Expected number of X_{17} in signal peak

$$N_{X17} = N_{ALL} \cdot f_{X17}$$


$$N_{X17} = 2758 \cdot 26.28\% = 725$$

General motivation – ATOMKI results (X_{17} boson)



The ATOMKI group observed an excess of e^+e^- pairs emitted at large relative angle in the nuclear reactions:




Expected ${}^8\text{Be}$ Transition



Hypothetical



Atomki Experiment



ATOMKI Exp. ${}^8\text{Be}$: anomalies in the internal pair creation of isovector (17.64 MeV, $I=1$) and isoscalar (18.15 MeV, $I=0$) magnetic dipole M1 transitions in ${}^8\text{Be}$



Light η , ω , f_1 mesons studies using **HADES**

- **Searches for Axion-Like-Particles $\eta \rightarrow \pi^+ \pi^- e^+ e^-$ Decay**
(this presentation)
- Exclusive and inclusive production of η and ω in proton-proton collisions (production mechanism at intermediate energies, cross section extraction, angular distributions).
- Form Factor extraction for the $\eta \rightarrow \gamma e^+ e^-$ and $\omega \rightarrow \pi^0 e^+ e^-$.
- Studies of the f_1 meson production in proton-proton (exclusive cross section extraction).
- Studies of symmetries C and CP using η decays:
 $\eta \rightarrow \pi^+ \pi^- e^+ e^-$, $\eta \rightarrow \pi^0 e^+ e^-$, $\eta \rightarrow \pi^+ \pi^- \pi^0$.

Marcin Zielinski
Krzysztof Prościński
(PhD Student)

Iza Ciepał
Szymon Treliński
(PhD Student)

Next analysis