# What is hiding at 750 GeV?

Veronica Sanz (Sussex)




KPS discovery, Daejeon

# Challenges


# Standard Model of Particle Physics

Predictive, successful paradigm being tested to higher and higher precision at the LHC





Based on QFT, symmetries (global/gauge) and consistent ways to break them
Foundation from which we develop theories beyond the SM



# Example of unified framework: Supersymmetry

Unifies concept of bosons and fermions

Light scalar bosons

Candidates for Dark Matter

Unification of strong/EM/weak forces


Matter/Antimatter asymmetry

Component of Quantum Gravity


New mechanisms
Inflation, Neutrinos and Dark Energy

The discovery of SUSY at LHC first step to understand many aspects of Nature

#### Nature, meanwhile



#### Nature, meanwhile



# The diphoton excess characteristics

(Theory) Non-trivial (Experiment) extremely clean

#### What is it?

An excess in a channel with two photons at an invariant mass of about 750 GeV

scalar, e.g. more Higgses tensor, e.g. spin-two graviton off-shell spin-one? (ask Felix)

What we knew before Dec 2015
Run 1: CMS already a (less significant) excess,
ATLAS did not show above 600 GeV

Dec 2015 excess in both ATLAS and CMS Run2 data


#### Moriond 2016

ATLAS and CMS results for s=0 & 2 narrow and wide

ATLAS analysis note public

CMS update including improvements in mass resolution and 0T data-set

## By eye



## Significance


ex. interpreted as a gluon-fusion narrow scalar (similar results for spin-two)

3.4 CMS Preliminary 3.3 fb<sup>-1</sup> (13 TeV) + 19.7 fb<sup>-1</sup> (8 TeV) മ് 1σ 10<sup>-1</sup> 2 σ 10<sup>-2</sup>  $\frac{\Gamma}{m} = 1.4 \times 10^{-4} \text{ J} = 0$ 3 σ 10<sup>-3</sup> Combined ----- 8TeV ---- 13TeV 10-4  $8 \times 10^{2}$ 5×10<sup>2</sup>  $6 \times 10^{2}$  $7 \times 10^{2}$ ms (GeV)

**CMS** 

**ATLAS** 


3.6



(remember LEE should be taken only once)

#### Production

Kick from 8 to 13 TeV from non-valence quarks or gluons




sizeable cross section & narrow resonance indicates gluon-initiated


but other productions, incl diphoton still an option

where are the photons? EBEB vs EBEE
Initially (Dec), it looked as if kinematics were funny

#### **CMS**




but s=0 and 2 are not so different




Han, Lee, Park, VS.

where are the photons? EBEB vs EBEE
Initially (Dec), it looked as if kinematics were funny


#### **CMS**





but we didn't have ATLAS to compare with


where are the photons? EBEB vs EBEE
Initially (Dec), it looked as if kinematics were funny



but we didn't have ATLAS to compare with

Is this signal coming along other objects?

#### 1. It doesn't recoil (much)



Is this signal coming along other objects?


2. No electrons or muons

e.g. from ATLAS analysis

"In addition, no electron or muon candidates have been found, with  $p_T > 10$  GeV and  $|\eta| < 2$ . (electrons) or 2.7 (muons) in the events with invariant masses between 700 GeV and 840 GeV.

Is this signal coming along other objects?


#### 3. No high-pT jets



jet anti-kT 0.4 pT> 25, eta< 4.4


Is this signal coming along other objects?


#### 4. No MET



#### Kinematics Narrow or wide?


#### CMS prefers narrow





#### **ATLAS**


slight preference wide (0.3 sigma)




overall no preference for wide

## Signal strength

# compatibility? Run1 vs Run 2 and CMS vs ATLAS





theorists combination in Dec

 $6.2 \pm 1.0 \text{ (fb)} \text{ (local)}$ 

#### Other final states

light Higgs into diphotons is not like the 750 GeV

Higgs below the threshold of WW, ZZ suppressed BRs

A heavy resonance in two photons? it couples to SM gauge interactions we expect WW, ZZ and Zgamma (and hh)

#### Model-independent prediction:

diphotons means there must be at least one non-zero BR(Z-gamma) and/or BR(ZZ)

$$g_{\gamma\gamma} = c_1 \alpha_1 c_W^2 + c_2 \alpha_2 s_W^2 \longrightarrow$$


$$g_{z\gamma} = (c_1 \alpha_1 - c_2 \alpha_2) s_{2W}$$
$$g_{zz} = c_1 \alpha_1 s_W^2 + c_2 \alpha_2 c_W^2$$

non-zero coupling to photons,

No, VS, Setford. 1512.0


coupling to ZZ and/or Zphoton





#### spin-0 vs spin-2 both compatible

Data (600 GeV< m', < 700 GeV) Data (700 GeV< m, < 840 GeV)



**LAS** Preliminary


= 13 TeV, 3.2 fb<sup>-1</sup>

# Models for the diphoton

Many papers written (~290 today)
Some model-independent,
most model-building

#### What is it, if anything?

maybe a scalar coupled to vector-like fermions?



SUSY, Composite scalar?

likely to be seen it in other channels with vector bosons

WW, ZZ, Z-photon

compatible with diphoton first

Spin

J=0

A new scalar

Would this be the end of anthropics?

Spin

J=0 A new scalar

Hooray SUSY!?

MSSM or NMSSM
will not do
compatibility with other
searches, dof,
perturbativity and tuning

non-minimal or threshold effects

## Spin

J=0 A new scalar

#### Hooray SUSY!?

MSSM or NMSSM
will not do
compatibility with other
searches, dof,
perturbativity and tuning

non-minimal or threshold effects

#### Composite dynamics?

glueball of new strong force or a pseudo-Goldstone boson

link to Composite Higgs

e.g.


see-saw composite Higgs Dark Matter, Baryogenesis

No, VS, Setford.1512.05700

# Spin J=2

# A kind of massive graviton or glueball of new strong force

#### Important hurdle is EWPTs



Spin
J=2

A kind of massive graviton or glueball of new strong force

Important hurdle is EWPTs

Experimental interpretations neglect this problem, theorists use AdS/CFT to find models

recent progress

1603.06980, 1603.08250


& in composite Higgs


Dillon, VS. 1603.09550

with DM: talk by HM Lee

#### Conclusions

- Two excesses at roughly 3.5 sigma on same location in mass at 750 GeV and cross section 5 fb LEE reduces each to about 2 sigma (double-counting)
- Width and spin still TBD. Excess doesn't come with high-pT objects. Most compatible with gluon-fusion
- Models of spin-zero: composite scalars or nonstandard SUSY
- Models of spin-two: non-standard AdS/CFT techniques required





Whatever is hiding, making sense of naturalness, Dark Universe and model-building techniques is a challenge for theorists.

300 papers in ~ 4 months, we are up to it!