

# On a reinterpretation of the Higgs field in supersymmetry and a proposal for new quarks

Carlos Muñoz



Instituto de  
Física  
Teórica  
UAM-CSIC




Departamento de  
Física Teórica

Madrid, Spain

MultiDark

Multimessenger Approach  
for Dark Matter Detection



DSU 2017, Daejeon, July 10-14, 2017

Work in collaboration with

**Daniel E. López-Fogliani**

(CONICET, Univ. Buenos Aires, Argentina)

**arXiv: 1701.02652, PLB**

# The Higgs is intriguing

- the only elementary scalar in the spectrum of the **SM**
- introduces the hierarchy problem
- no three-fold replication, unlike the rest of the matter

In **SUSY** the presence of the Higgs(es) is more natural:

- scalar particles exist by construction
- the hierarchy problem can be solved
- models predict the Higgs mass  $\lesssim 140$  GeV

**BUT**

- NO explanation is given for the existence of only one family
- Their behaviour is very different from the rest of the matter

# Supersymmetry

- Higgses do not have a three-fold replication as the rest of the matter

$$L_i = \begin{pmatrix} \nu_i \\ e_i \end{pmatrix}, \quad e_i^c, \quad Q_i = \begin{pmatrix} u_i \\ d_i \end{pmatrix}, \quad d_i^c, \quad u_i^c \quad \begin{array}{l} \text{3 families of superfields in SUSY} \\ \text{sleptons + leptons} \quad \text{squarks + quarks} \end{array}$$

$Y = -1/2$

$$H_d = \begin{pmatrix} H_d^0 \\ H_d^- \end{pmatrix}, \quad H_u = \begin{pmatrix} H_u^+ \\ H_u^0 \end{pmatrix} \quad \begin{array}{l} \text{1+1 Higgs doublet superfields (vector-like) are needed} \\ \text{W} = Y_e H_d^T \epsilon L e^c + Y_d H_d^T \epsilon Q d^c - Y_u H_u^T \epsilon Q u^c \end{array}$$

$Y = -1/2 \quad Y = +1/2$

but since  $H_d$  and  $L$  have the same SM quantum numbers,  $Y = -1/2$

this might lead one to interpret the Higgs superfield  $H_d$  as a fourth family of lepton superfields

$$L_4 = \begin{pmatrix} \nu_4 \\ e_4 \end{pmatrix} = \begin{pmatrix} H_d^0 \\ H_d^- \end{pmatrix} = H_d$$

Unfortunately, one cannot interpret naturally the superfield  $H_u$  in a similar way, given that it is a doublet with no leptonic counterpart for its neutral component (...right-handed neutrinos)

$$L_4 = \begin{pmatrix} \nu_4 \\ e_4 \end{pmatrix} = \begin{pmatrix} H_d^0 \\ H_d^- \end{pmatrix} = H_d$$

- Actually, even  $H_d$  cannot be interpreted as a fourth family of leptons, since
  - there can be no mixing between particles and sparticles,

Higgses are NOT sleptons and Higgsinos are NOT leptons

thus their behaviour is very different, e.g.:

$$\begin{aligned} \langle H_u^0 \rangle &\neq 0, \quad \langle H_d^0 \rangle \neq 0, \\ \langle \tilde{v}_{eL} \rangle &= 0, \quad \langle \tilde{v}_{\mu L} \rangle = 0, \quad \langle \tilde{v}_{\tau L} \rangle = 0 \end{aligned}$$

This connection is **ONLY** possible if R-parity is violated



e.g. the Yukawa couplings of the **MSSM**


$$W = Y_{ij}^e H_d L_i e_j^c + Y_{ij}^d H_d Q_i d_j^c - Y_{ij}^u H_u Q_i u_j^c$$

have effectively a  $Z_2$  discrete symmetry (**R parity**)

$$R_p(\text{particle}) = +1$$

$$R_p(\text{sparticle}) = -1$$

i.e. sparticles must appear in pairs



because of these couplings, there are **mixing between sparticles**, e.g. gauginos & higgsinos

$$(\tilde{B}^0, \tilde{W}^0, \tilde{H}_1^0, \tilde{H}_2^0) \longrightarrow \mathcal{M} = \begin{pmatrix} M_1 & 0 & -\frac{g'\nu_1}{\sqrt{2}} & \frac{g'\nu_2}{\sqrt{2}} \\ 0 & M_2 & \frac{g\nu_1}{\sqrt{2}} & -\frac{g\nu_2}{\sqrt{2}} \\ -\frac{g'\nu_1}{\sqrt{2}} & \frac{g\nu_1}{\sqrt{2}} & 0 & -\mu \\ \frac{g'\nu_2}{\sqrt{2}} & -\frac{g\nu_2}{\sqrt{2}} & -\mu & 0 \end{pmatrix}$$

4 neutralinos:  $\tilde{\chi}_1^0 = N_{11}\tilde{B}^0 + N_{12}\tilde{W}^0 + N_{13}\tilde{H}_1^0 + N_{14}\tilde{H}_2^0$   
 2 charginos:  $(\tilde{W}^+, \tilde{H}^+)$

**BUT because of the different  $R_p$  quantum numbers, there can be no mixing between particles and sparticles**, e.g. between neutralinos and neutrinos, ...


# Supersymmetry with right-handed neutrinos

The gauge-invariant trilinear superpotential containing right-handed neutrinos:

Lopez-Fogliani, C. M., PRL 2006

$$W = Y_{ij}^e \boxed{H_d} L_i e_j^c + Y_{ij}^d \boxed{H_d} Q_i d_j^c - Y_{ij}^u H_u Q_i u_j^c - Y_{ij}^\nu H_u \boxed{L_i} \nu_j^c + \lambda_{ijk} \boxed{L_i} L_j e_k^c + \lambda'_{ijk} \boxed{L_i} Q_j d_k^c + \frac{1}{3} \kappa_{ijk} \nu_i^c \nu_j^c \nu_k^c + \lambda_i H_u \boxed{H_d} \nu_i^c.$$

violates explicitly  $R_p$




$$\langle H_u^0 \rangle, \langle H_d^0 \rangle, \langle \tilde{\nu}_{iL} \rangle, \langle \tilde{\nu}_{iR} \rangle \neq 0$$

Fields with the same color, electric charge and spin naturally mix, e.g.:

"Neutrinos"

$$(\tilde{B}^0, \tilde{W}^0, \tilde{H}_d, \tilde{H}_u, \nu_{R_i}, \nu_{L_i}),$$



Parameter which determines the violation of  $R_p$ :  $Y^\nu$

$Y^\nu \longrightarrow 0$  the  $\nu^c$  are no longer neutrinos, they are just ordinary singlets like the  $N$  of the NMSSM:  $N H_1 H_2 + NNN$ , and R-parity is conserved

# A simple re-interpretation of the spectrum

neutrinos/leptons

neutral Higgs

exist in Nature (SM)

SUSY predicts

neutrinos/leptons

neutral/charged Higgsinos

are mixed


+

+

sneutrinos/sleptons

neutral/charged Higgses

are mixed



in SUSY with  
neutrinos/RPV

There are only neutrinos/leptons (and quarks) and their scalar partners

$$L_i = \begin{pmatrix} \nu_i \\ e_i \end{pmatrix}, \quad \begin{matrix} e_i^c \\ \nu_i^c \end{matrix}, \quad Q_i = \begin{pmatrix} u_i \\ d_i \end{pmatrix}, \quad \begin{matrix} d_i^c \\ u_i^c \end{matrix} \quad \longrightarrow \quad L_i = \begin{pmatrix} \nu_i \\ e_i \end{pmatrix}, \quad \begin{matrix} e_i^c \\ \nu_i^c \end{matrix}, \quad Q_i = \begin{pmatrix} u_i \\ d_i \end{pmatrix}, \quad \begin{matrix} d_i^c \\ u_i^c \end{matrix}$$

$$H_d = \begin{pmatrix} H_d^0 \\ H_d^- \end{pmatrix}, \quad H_u = \begin{pmatrix} H_u^+ \\ H_u^0 \end{pmatrix} \quad \text{Higgses are vector-like leptons of a fourth family} \quad \longrightarrow \quad L_4 = \begin{pmatrix} \nu_4 \\ e_4 \end{pmatrix}, \quad L_4^c = \begin{pmatrix} e_4^c \\ \nu_4^c \end{pmatrix}.$$

In this framework, the first scalar particle discovered at the LHC is a sneutrino belonging to a 4<sup>th</sup>-family vector-like doublet representation



## Extra bonuses

$$\begin{aligned}
 W = & Y_{ij}^e H_d L_i e_j^c + Y_{ij}^d H_d Q_i d_j^c - Y_{ij}^u H_u Q_i u_j^c - Y_{ij}^\nu H_u L_i \nu_j^c \\
 & + \lambda_{ijk} L_i L_j e_k^c + \lambda'_{ijk} L_i Q_j d_k^c + \frac{1}{3} \kappa_{ijk} \nu_i^c \nu_j^c \nu_k^c + \lambda_i H_u H_d \nu_i^c .
 \end{aligned}$$

When the right sneutrinos acquire VEVs of order the EW scale, an effective  $\mu$ -term from  $\nu$  is generated ( $\mu\nu$ SSM)

Lopez-Fogliani, C. M., PRL 2006

-producing higgsino masses beyond the experimental bounds  $\mu \geq 100$  GeV

**solving the  $\mu$  problem:** What is the origin of  $\mu$ , and why is so small  $\ll M_{\text{Planck}}$

-as well as effective Majorana masses for neutrinos: EW scale seesaw

$$m_\nu \sim m_D^2/M_M = (\mathbf{Y}_\nu v_u)^2/(k v_R) \sim (10^{-6} 10^2)^2/10^3 = 10^{-11} \text{ GeV} = 10^{-2} \text{ eV}$$



Like the electron Yukawa

**solving the  $\nu$  problem:** How to accommodate the neutrino data

## Extra bonuses


- Since R parity is violated, SUSY particles can decay to standard model particles, there is no missing energy as a special signal, and the bounds become significantly weaker
- Interesting LHC phenomenology because of R-parity violation
  - Novel signals with displaced vertices, multilepton final states, multijets  
...

E.g.:

# Left Sneutrino LSP

$$\tilde{\nu}_{iL} \quad i=e,\mu,\tau$$

Ghosh, Lara, Lopez-Fogliani, C.M., Ruiz de Austri, arXiv:1707.02471



$$Y_v \sim 10^{-6}$$



$$45 \lesssim m_{\tilde{\nu}_L} \lesssim 300 \text{ GeV}$$

in order to observe its production at the LHC

$$m_{\tilde{\nu}_L} \sim 45 - 100 \text{ GeV} \quad \text{have decay lengths} \sim \text{mm}$$

**DISPLACED**


## Proposal for new quarks

$$L_i = \begin{pmatrix} \nu_i \\ e_i \end{pmatrix}, \quad \begin{matrix} e_i^c \\ \nu_i^c \end{matrix}, \quad Q_i = \begin{pmatrix} u_i \\ d_i \end{pmatrix}, \quad \begin{matrix} d_i^c \\ u_i^c \end{matrix}, \quad \text{For the first 3 families, each lepton representation has its quark counterpart}$$

$$L_4 = \begin{pmatrix} \nu_4 \\ e_4 \end{pmatrix}_{Y=-1/2}, \quad L_4^c = \begin{pmatrix} e_4^c \\ \nu_4^c \end{pmatrix}_{Y=+1/2}, \quad Q_4 = \begin{pmatrix} u_4 \\ d_4 \end{pmatrix}_{Y=+1/6}, \quad Q_4^c = \begin{pmatrix} d_4^c \\ u_4^c \end{pmatrix}_{Y=-1/6}$$

In analogy, we add to the 4<sup>th</sup> family a **vector-like quark doublet representation** as counterpart of the vector-like lepton/Higgs doublet representation

$$\begin{aligned} W = & Y_{ij}^e H_d L_i e_j^c + Y_{ij}^d H_d Q_i d_j^c - Y_{ij}^u H_u Q_i u_j^c - Y_{ij}^\nu H_u L_i \nu_j^c \\ & + \lambda_{ijk} L_i L_j e_k^c + \lambda'_{ijk} L_i Q_j d_k^c + \frac{1}{3} \kappa_{ijk} \nu_i^c \nu_j^c \nu_k^c + \lambda_i H_u H_d \nu_i^c \\ & + \lambda'_{i4k} L_i Q_4 d_k^c + Y_{4k}^d H_d Q_4 d_k^c - Y_{4k}^u H_u Q_4 u_k^c + Y_{j4k}^Q Q_j Q_4^c \nu_k^c + Y_{44k}^Q Q_4 Q_4^c \nu_k^c \end{aligned}$$

Mass term for the new quarks 

**We note in passing...**

**Strings?**

The presence of extra vector-like matter is a common situation in string constructions:  
Orbifolds, Branes,...

Font, Ibáñez, Nilles, Quevedo, 88; Font, Ibáñez, Quevedo, Sierra, 90  
Casas, Katehou, C.M., 87; Casas, C.M., 88  
Cvetic, Shiu, Uranga, 01

...

# Novel signatures for vector-like quarks

**arXiv: 1705.02526, JHEP**

Work in collaboration with


**J. A. Aguilar-Saavedra**  
(Univ. Granada, Spain)

**D. E. López-Fogliani**  
(CONICET, Univ. Buenos Aires, Argentina)



# Detection of heavy vector-like quarks (T B) at the LHC

Many searches for heavy quark pair or single production focus on the standard decay modes:







New BRs are possible in models with non-minimal scalar sectors, modifying the analyses:

$$\begin{array}{lll} T \rightarrow H^+ b & T \rightarrow H_k^0 t & T \rightarrow P_k^0 t \\ B \rightarrow H^- t & B \rightarrow H_k^0 b & B \rightarrow P_k^0 b \end{array} \left. \vphantom{\begin{array}{lll} T \rightarrow H^+ b \\ B \rightarrow H^- t \end{array}} \right\} \begin{array}{l} k=1 \longrightarrow \text{MSSM like} \\ k=1,2 \longrightarrow \mu\nu\text{SSM } \tilde{V}_R \end{array}$$

New signals can be produced:

$$\begin{array}{ll} T \rightarrow t \bar{t} t & T \rightarrow h^0 h^0 t \\ B \rightarrow t \bar{t} b & B \rightarrow h^0 h^0 b \end{array}$$

# Detection of heavy vector-like quarks (T B) at the LHC



$$T \rightarrow t \bar{b} b$$

$$B \rightarrow t \bar{t} b$$

$$B \rightarrow h^0 h^0 b$$

# The lower limits on the (T,B) quark masses are less stringent

$$\mathbf{T} \rightarrow W^+ b \quad \mathbf{T} \rightarrow Z t \quad \mathbf{T} \rightarrow h^0 t \quad \text{Br}(W) + \text{Br}(Z) + \text{Br}(h^0) = \rho = 1$$

New decay modes  $\longrightarrow$   $\text{Br}(W) + \text{Br}(Z) + \text{Br}(h^0) = \rho < 1$




Figure 6: Lower limits on the  $T$  quark mass for several values of  $\rho$  from a recast of the limits of the heavy quark search in ATLAS-CONF-2016-101

# Conclusions

SUSY with right-handed neutrinos naturally produces R-parity violation

- Interesting theoretical advantages:

- solves the  $\mu$  problem
- solves the  $v$  problem
- reinterpretation of the Higgs(es) as a “4<sup>th</sup> family” of lepton superfields

- Interesting LHC phenomenology:

- Novel signals with displaced vertices,  
multilepton final states,  
multijets  
diphoton + leptons  
diphoton + missing energy

THE END

new vector-like quarks:

$$\mathbf{T} \longrightarrow t \bar{t} t \quad \mathbf{T} \longrightarrow h^0 h^0 t \quad \mathbf{B} \longrightarrow t \bar{t} b \quad \mathbf{B} \longrightarrow h^0 h^0 b$$



## DSU 2005

SURVIVORS OF THE WORKSHOP PREPARED TO ENTER IN THE  
DARK SIDE OF THE UNIVERSE

