

Nonthermal heavy dark matter from a first-order phase transition

Hyun Min Lee
Chung-Ang University



Ref. G. Giudice, HML, A. Pomarol, B. Shakya, JHEP12 (2024) 190.

Focus workshop on cosmological phase transitions
IBS CTPU, Daejeon, Nov 22, 2025

Outline







- Introduction
- Dark matter from bubble collisions
- Vector dark matter and gauge choices
- Conclusions

Introduction

Origin of dark matter




-1-

We don't have a clue to the nature of 27% dark matter and 68% dark energy.



Bounds on WIMP

-2-



Solar (^8B) neutrinos ($2.6-2.7\sigma$),
e.g. 2408.02877

- WIMP-nucleon cross section strongly constrained by direct detection, such as XENON experiments.
- New DM: Axion-like DM, leptophilic DM, resonant DM, etc.


Dark matter production

-3-

DM in thermal equilibrium

- Freeze-out mechanism

DM annihilation to thermal bath




$$\Omega_{\text{DM}} \sim \frac{1 \text{ pb}}{\langle \sigma v \rangle}$$

DM out of equilibrium

- Freeze-in mechanism

Tiny DM-SM interaction




$$\langle \sigma v \rangle = \sigma_0 (T/m_{\text{DM}})^n, \quad T > m_{\text{DM}}$$

$$\Omega_{\text{DM}} \propto \sigma_0 (T_{\text{RH}})^{n+1}, \quad n > -1$$

$n < -1 : T_{\text{RH}}\text{-independent}$

- DM self-interactions

$N(\geq 3) \rightarrow 2$ annihilation but
small DM-SM interaction




$$\Omega_{\text{DM}} \sim \frac{1 \text{ pb}}{(\alpha_{\text{self}} \sigma_{\text{self}} / (m_{\text{DM}} M_P))^{1/2}}$$

$$\sigma_{\text{self}} = \frac{\alpha_{\text{self}}^2}{m_{\text{DM}}^2} \sim 1 \text{ cm}^2/\text{g}$$

- Misalignment (boson DM)

DM initial potential density

Ultra light DM $m_a \sim H \sim \frac{T_{\text{osc}}^2}{M_P}$





$$\Omega_a h^2 \simeq 0.12 \left(\frac{f_a}{9 \times 10^{11} \text{ GeV}} \right)^{1.165} \left(\frac{a_{\text{ini}}}{f_a} \right)^2$$

Bounds on DM masses

-4-


- WIMP/SIMP: Unitarity bounds



$$(\sigma v)_{\text{th}} \simeq 3 \times 10^{-26} \text{ cm}^3/\text{s} < \frac{16\pi}{m_{\text{DM}}^2 v} \longrightarrow m_{\text{DM}} \lesssim 100 \text{ TeV}$$


$$\frac{\alpha_{\text{self}} \sigma_{\text{self}}}{m_{\text{DM}}} \simeq 10^{-2} \text{ cm}^2/\text{g} < \frac{16\pi \alpha_{\text{self}}}{m_{\text{DM}}^3 v^2} \longrightarrow m_{\text{DM}} \lesssim 1 \text{ GeV}$$

- FIMP: Temperature bound




$$\langle \sigma v \rangle \propto (\sigma v) \Big|_{T=m_{\text{DM}}} \times e^{-m_{\text{DM}}/T} \longrightarrow m_{\text{DM}} \lesssim T_{\text{RH}}$$

- DM mass beyond unitarity or temperature bound?
- Evolution history (reheating, phase transitions, axion rotation)?

Bounds on DM masses

- Reheating period or low reheating temperature

-5-






$$T_{RH} \lesssim m_{DM} \lesssim T_{max}, \quad \langle \sigma v \rangle \propto (\sigma v) \Big|_{T=m_{DM}} \times e^{-m_{DM}/T}$$

“Larger cross section”

Stronger couplings than FIMP (“Strong freeze-in”)

[Cosme et al (2023); Arcadi et al (2024);
Bernal et al (2024); Sanz, HML, Park (2024)]







QCD axion rotation

-6-

$$\epsilon_{\theta, \text{end}} = 1$$



[Co, Hall, Harigaya (2019); HML, A. Menkara, M-J, Seung, J-H, Song (2023, 2024)]

$$\epsilon_\theta \equiv \frac{M_P^2}{2V_E^2} \left(\frac{\partial V_E}{\partial \theta} \right)^2 \longrightarrow Y_\theta = \frac{n_\theta(T_{\text{RH}})}{s(T_{\text{RH}})} \longrightarrow \Omega_a h^2 = 0.12 \left(\frac{10^9 \text{ GeV}}{f_a} \right) \left(\frac{Y_\theta}{40} \right)$$



Axion velocity from inflation $n_\theta = f_a^2 \dot{\theta}$

Smaller axion decay or
larger axion mass

Phase transition and DM

- DM can receive mass via 1st-order phase transitions.


-7-



Inside bubbles:
true vacuum

Outside bubbles:
false vacuum

[M. Hindmarsh et al, 2020]



Outside bubbles Inside bubbles

[M. Baker et al ; D. Chway et al, 2019]

DM abundance is filtered for a non-relativistic bubble wall, $v_w \ll 1$.


DM interactions of order one:

$$\Omega_{\text{DM}} h^2 \sim 0.1 \left(\frac{T_n}{1 \text{ TeV}} \right) \left(\frac{x}{30} \right)^{5/2} e^{-(x-30)}, \quad x = \frac{m_{\text{DM}}}{T_n}$$

$m_{\text{DM}} \sim 30 T_n$, DM mass beyond T.

Bubble nucleation

-8-



$T \gg R_0^{-1}$: $O(3)$ symmetric bubble

Bounce action: $S_4 = \int d\tau d^3x \mathcal{L}_E = \frac{S_3}{T},$

$$S_3 = 4\pi \int_0^\infty s^2 ds \left[\frac{1}{2} \left(\frac{d\phi}{ds} \right)^2 + V_T(\phi) \right], \quad s = |\vec{x}|$$

Bubble profile: $\frac{d^2\phi}{ds^2} + \frac{2}{s} \frac{d\phi}{ds} - V'_T(\phi) = 0,$ w/ b.c. $\frac{d\phi}{dr} = 0,$ at $r = 0,$
 $\phi = 0,$ at $r = \infty.$

Thin-wall approximation: $l_0 \ll R_0$


$$S_3 \simeq 4\pi\sigma R^2 - \frac{4\pi}{3}\Delta V R^3 = \frac{16\pi}{3} \frac{\sigma^3}{(\Delta V)^2} \longrightarrow R_0 = \frac{2\sigma}{\Delta V} \quad \text{“critical radius”}$$

Bubble nucleation rate & inverse duration time:

$$\Gamma_n \sim T_n^4 e^{-S_4(T_n)} \sim H(T_n), \quad \frac{\beta}{H_*} = T \frac{dS_4}{dT} \Big|_{T_*} - 4 \gtrsim 1$$

Running bubbles

-9-



Bubble dynamics:

$$\mathcal{L} = -2\pi\sigma R^2 \sqrt{1 - \dot{R}^2} + \frac{4\pi}{3} R^3 p,$$

$$p = \Delta V - \Delta p_{\text{LO}} - \gamma \Delta p_{\text{NLO}}$$

→ e.o.m: $\frac{d\gamma}{dR} + \frac{2\gamma}{R} = \frac{p}{\sigma}$


1) Difference in FT potential:

$$\Delta p_{\text{LO}} = \Delta V_{\text{FT}} = \frac{1}{24} (\Delta m^2) T^2$$

[Boedeker, Moore, 2009]

See also I. Baldes, 2024.


$$\gamma_{\text{max}} = \frac{2R_{\text{max}}}{3R_0}, \quad R_{\text{max}} \sim v_w \beta^{-1}$$



$$\Delta V = c_V v_\phi^4, \quad T \sim v_\phi \quad \longrightarrow \quad \gamma_{\text{max}} \sim \frac{1}{(\beta/H_*)} \frac{M_P}{v_\phi} \gg 1$$

2) Pressure due to radiation of light particles:


[Boedeker, Moore, 2017; Gouttenoire et al, 2021] $\gamma \Delta p_{\text{NLO}} = \gamma g^2 \Delta m_V T^3$




$$p = 0 \quad \longrightarrow \quad \gamma_{\text{eq}} = \frac{\Delta V - \Delta p_{\text{LO}}}{\Delta p_{\text{NLO}}} \simeq \frac{\Delta V}{g^3 v_\phi T^3} \sim \frac{c_V}{g^3} \gg 1, \quad g \ll 1$$

Bubble collisions: DM & GW

-10-



[G. Giudice et al, 2024]



[K. Schmitz, 2020]

DM density:
$$\Omega_\chi h^2 \approx 0.1 \left(\frac{\beta/H}{10} \right) \left(\frac{\alpha}{(1+\alpha)g_*c_V} \right)^{1/4} \frac{m_\chi v_\phi}{(2.5 \text{ TeV})^2} \int_{p_{\min}^2}^{p_{\max}^2} \frac{dp^2}{p^4} \text{Im}[\tilde{\Gamma}^{(2)}(p^2)]$$

GW spectrum:
$$\Omega_{\text{coll}}(f) h^2 = 1.66 \times 10^{-5} \left(\frac{H_*}{\beta} \right)^2 \frac{\kappa_{\text{coll}}^2 \alpha^2}{(1+\alpha)^2} \left(\frac{100}{g_*(T_*)} \right)^{1/3} \frac{v_w^3}{1+2.4v_w^2} \cdot \frac{(f/f_{\text{coll}})^{2.8}}{1+2.8(f/f_{\text{coll}})^{3.8}}$$

$$\alpha = \frac{\Delta V}{\rho_R}, \quad v_\phi \sim T$$

Experiment	$f_{\text{optimal}}/\text{Hz}$	v_ϕ/GeV	m_{DM}/GeV
Pulsar Timing Arrays (PTAs) [107]	10^{-8}	0.1	$10^{13} - 10^{16}$
LISA [108]	0.001	10^4	$10^6 - 10^{15}$
BBO [109], DECIGO [110]	0.1	10^6	$10^5 - 10^{13}$
Einstein Telescope (ET) [111], Cosmic Explorer (CE) [112]	10	10^8	$10^6 - 10^{10}$

Dark matter from bubble collisions

Green function formalism

Probability for particle production: $\mathcal{P} = 2 \text{Im}(\Gamma[\phi])$ -11-

$\Gamma[\phi]$, Effective action or generating function of 1PI Green functions

Quadratic in ϕ : $\Gamma[\phi] = \sum_{n=2}^{\infty} \frac{1}{n!} \int d^4x_1 \dots d^4x_n \Gamma^{(n)}(x_1, \dots, x_n) \phi(x_1) \dots \phi(x_n)$.

$$\longrightarrow \mathcal{P} = \int \frac{d^4p}{(2\pi)^4} |\tilde{\phi}(p)|^2 \text{Im}(\tilde{\Gamma}^{(2)}(p^2)), \quad \tilde{\phi}(p) = \int d^4x \phi(x) e^{ip \cdot x}, \text{ etc.}$$

Bubble profile in momentum space: $\tilde{\phi}(p) = (2\pi)^2 \delta(p_x) \delta(p_y) \tilde{\phi}(p_z, \omega)$

Number density of particles per area: [R. Watkins, L.M. Widrow, 1992]

$$\frac{N}{A} = 2 \int \frac{dp_z d\omega}{(2\pi)^2} |\tilde{\phi}(p_z, \omega)|^2 \text{Im}[\tilde{\Gamma}^{(2)}(\omega^2 - p_z^2)]$$

Background dynamics



Particle physics info.

Bubble collisions: elastic

-12-

Almost degenerate minima: elastic bubble collision

=> scalar waves, particle production



[A. Falkowski, J.M. No, 2012]

Thin wall: $\phi(z, t) = \begin{cases} 0 & , \quad z^2 < v_w^2 t^2, \\ v_\phi & , \quad z^2 > v_w^2 t^2. \end{cases} \longrightarrow \tilde{\phi}(p_z, \omega) = \frac{4v_w v_\phi}{\omega^2 - v_w^2 p_z^2}$

Running bubbles, $v_w \simeq 1 \longrightarrow \tilde{\phi}(p_z, \omega) \simeq \frac{4v_\phi}{p^2}$, IR dominant

Thick wall: $\phi(z, t) = \frac{v_\phi}{2} \left(2 + \tanh\left(\frac{z - |t|}{l_w}\right) - \tanh\left(\frac{z + |t|}{l_w}\right) \right)$, $l_w = \frac{l_0}{\gamma_w}$.

$\longrightarrow \tilde{\phi}(p, \omega) \simeq \frac{2\pi l_w v_\phi \omega}{\sinh(\frac{1}{2}\pi l_w \omega)} \frac{1}{p^2} \propto e^{-\pi l_w \omega/2}, \quad \omega \gg 1/l_w.$


Bubble collisions: elastic

-13-

Produced particles on surface diffuse over the bubble volume.

$$n = \frac{N}{V} = \frac{3}{4\pi^2 R_*} \int_{p_{\min}^2}^{p_{\max}^2} dp^2 f(p^2) \text{Im} \left[\tilde{\Gamma}^{(2)}(p^2) \right],$$

$$p_{\max} = 2/l_w, \quad p_{\min} = \min(2m, (2R_*)^{-1})$$



“Efficiency” factor for particle production

$$f(p^2) = \frac{1}{4} \int_{p^2}^{\infty} \frac{d\Psi}{\sqrt{\Psi^2 - p^4}} |\tilde{\phi}(p_z, \omega)|^2, \quad p^2 = \omega^2 - p_z^2$$



Elastic bubbles: $\tilde{\phi}(p, \omega) \simeq \frac{2\pi l_w v_\phi \omega}{\sinh(\frac{1}{2}\pi l_w \omega)} \frac{1}{p^2} \propto e^{-\pi l_w \omega/2}, \quad \omega \gg 1/l_w.$

$$\omega \sim m \sim \frac{\gamma_w}{l_0} \sim \gamma_w v_\phi \gg T \quad \Rightarrow \text{unsuppressed production}$$

Bubble collisions: inelastic

-14-

Non-degenerate minima: inelastic bubble collision
 $\Rightarrow$ scalar waves, mainly



[A. Falkowski, J.M. No, 2012]

Equation for bubble profile: $(\partial_t^2 - \partial_z^2)\phi = -\frac{\partial V}{\partial \phi}$, $t > 0$: $V(\phi) \simeq \frac{1}{2}m_\phi^2(\phi - v_\phi)^2$,



$$\longrightarrow \phi(z, t) = v_\phi + v_\phi l_w \int_0^\infty dp_z \frac{p_z}{\sqrt{p_z^2 + m_\phi^2}} \frac{\cos(p_z z)}{\sinh(\frac{1}{2}\pi l_w p_z)} \sin(\sqrt{p_z^2 + m_\phi^2} t), \quad t > 0.$$

$$\longrightarrow \tilde{\phi}(p_z, \omega) = \frac{\pi l_w v_\phi p_z}{\sinh(\frac{1}{2}\pi l_w p_z)} \left(\frac{1}{\omega^2 - p_z^2} - \frac{1}{\omega^2 - p_z^2 - m_h^2} \right),$$

$$m_{\text{DM}} \gg m_h : \quad f(p^2) \simeq f(p^2) \Big|_{\text{elastic}}$$

Elastic vs inelastic collisions

-15-



Perfectly Elastic:
$$f_{\text{PE}}(p^2) = \frac{16v_\phi^2}{p^4} \text{Log} \left[\frac{2(1/l_w)^2 - p^2 + 2(1/l_w)\sqrt{(1/l_w)^2 - p^2}}{p^2} \right]$$

[A. Falkowski, J.M. No, 2012]

Elastic collision describes well beyond the resonance.

2-body vs 3-body decays

-16-

$\tilde{\Gamma}^{(2)}(p^2)$: 2-point 1PI Green function (or self-energy)


Optical theorem: $\text{Im}[\tilde{\Gamma}^{(2)}(p^2)] = \frac{1}{2} \sum_k \int d\Pi_k |\mathcal{M}(\phi_p^* \rightarrow \underline{k})|^2$
 sum over all decays of off-shell scalar

Scalar self-interaction: $\frac{\lambda_\phi}{4!} \phi^4 \longrightarrow \frac{\lambda_\phi}{3!} v_\phi \bar{\phi}^3 + \frac{\lambda_\phi}{4!} \bar{\phi}^4$


2-body decay

3-body decay

[G. Giudice et al, 2024]



$$\text{Im}[\tilde{\Gamma}^{(2)}(p^2)]_{\phi_p^* \rightarrow \phi\phi} = \frac{\lambda_\phi^2 v_\phi^2}{8\pi} (1 - 4m_\phi^2/p^2) \Theta(p - 2m_\phi)$$



$$\text{Im}[\tilde{\Gamma}^{(2)}(p^2)]_{\phi_p^* \rightarrow 3\phi} = \frac{\lambda_\phi^2 p^2}{3072 \pi^3} (1 - 9m_\phi^2/p^2) \Theta(p - 3m_\phi)$$

$p^2 \gg v_\phi^2$: highly off-shell $\Rightarrow$ 3-body process dominates!

Scalar vs fermion DM



-17-

Assume that DM mass receives minor corrections from SSB.

Scalar DM χ_s

[G. Giudice et al, 2024]

$$\mathcal{L}_{\chi_s} = -\frac{\lambda_s}{4}\phi^2\chi_s^2 \longrightarrow \Delta m_{\chi_s}^2 = \frac{1}{2}\lambda_s v_\phi^2 \lesssim m_{\chi_s}, \quad \mathcal{L}_{\text{int}} = -\frac{\lambda_s}{2}v_\phi\bar{\phi}\chi_s^2$$



$p^2 \sim m_{\chi_s}^2 \gg v_\phi^2$: 3-body process dominant for SDM production.

Fermion DM χ_f

$$\mathcal{L}_{\chi_f} = -y_f\phi\bar{\chi}_f\chi_f \longrightarrow \Delta m_{\chi_f} = y_f v_\phi \lesssim m_{\chi_f}$$

Fermion DM self-energy:

2-body process is sufficient.

$$\text{Im}[\tilde{\Gamma}^{(2)}(p^2)]_{\phi_p^* \rightarrow \chi_f \bar{\chi}_f} = \frac{y_f^2}{8\pi} p^2 (1 - 4m_{\chi_f}^2/p^2)^{3/2} \Theta(p^2 - 4m_{\chi_f}^2) \sim p^2$$


Scalar DM relic density

-18-

Scalar DM can be produced through 2- & 3-body decays.

Scalar DM χ_s

$$\frac{\lambda_s}{4} \phi^2 \chi_s^2 \longrightarrow \frac{\lambda_s}{4} (v_\phi + \bar{\phi})^2 \chi_s^2 \rightarrow \frac{\lambda_s}{2} v_\phi \bar{\phi} \chi_s^2$$



DM relic density:

[G. Giudice et al, 2024]

$$\Omega_\chi h^2 \approx 0.1 \frac{\beta/H}{10} \left(\frac{\alpha}{(1+\alpha)g_*c_V} \right)^{1/4} \frac{\lambda_s^2 m_{\chi_s} v_\phi}{(24 \text{ TeV})^2} \left[\underbrace{\frac{v_\phi^2}{m_{\chi_s}^2}}_{\text{2-body}} + \underbrace{\frac{1}{16\pi^2} \ln \left(\frac{2\gamma_w/l_{w0}}{(2m_{\chi_s} + m_\phi)} \right)}_{\text{3-body}} \right]$$

$$\alpha = \frac{\Delta V}{\rho_R}, \quad \Delta V = c_V v_\phi^4$$

$m_{\chi_s} \gg v_\phi \sim T \longrightarrow$ dominant 3-body process!

Wall-plasma interactions

[Azakov, Vanvlasselaer, 2020; + W. Yin, 2021]

A wall scalar with boost can split into DM.

1. kinematic condition: $p_\phi = \gamma T_* > 2m_{\chi_s}$

2. non-adiabatic condition: $(\Delta p_z)l_0 \lesssim 1$, $\Delta p_z = \frac{m_{\chi_s}^2}{2p_\phi}$

$$\gamma T_* \gtrsim m_{\chi_s}^2 l_0 \sim \frac{m_{\chi_s}^2}{v_\phi} : \quad m_{\chi_s} \gg v_\phi \longrightarrow \gamma T_* \gg m_{\chi_s}$$

Heavy DM

much stronger than kinematic condition.


Thus, wall-plasma effects suppressed for mildly boosted wall.

Extra relic abundance:

$$\Omega_\chi h^2 \approx \frac{1.35 \times 10^5 \lambda_s^2}{g_*} \frac{v_\phi^2}{m_\chi^2 \text{ GeV}} \left(\frac{T_n}{T_*} \right)^3$$



[G. Giudice et al, 2024]

[See also J. Cembranos et al, 2024.]



Bubble collisions for DM

-20-



$c_V = 0.1, \beta/H = 10, \alpha = 1, R_0 = 10/v_\phi, l_{w0} = 1/v_\phi$ [G. Giudice et al, 2024]

DM density:
$$\Omega_\chi h^2 \approx 0.1 \left(\frac{\beta/H}{10} \right) \left(\frac{\alpha}{(1+\alpha)g_*c_V} \right)^{1/4} \frac{m_\chi v_\phi}{(2.5 \text{ TeV})^2} \int_{p_{\min}^2}^{p_{\max}^2} \frac{dp^2}{p^4} \text{Im}[\tilde{\Gamma}^{(2)}(p^2)]$$


Heavy dark matter: $m_{\chi_s} \gg v_\phi \sim T_*$ $\alpha = \frac{\Delta V}{\rho_R}, v_\phi \sim T$

→ 3-body decay process from bubble collision.

Scalar dark matter

-21-

Lyman- α
bounds
on hot DM



Portal for scalar DM:

$$\mathcal{L}_{\chi_s} = \frac{\lambda_s}{4} \phi^2 \chi_s^2$$

$$m_{\chi_s} < R_*^{-1}$$

Multiple bubbles
in the Fourier
analysis needed.

$$E_{\chi_s} \gtrsim R_*^3 \Delta V$$

$$c_V = 0.1, \beta/H = 10, \alpha = 1, R_0 = 10/v_\phi, l_{w0} = 1/v_\phi$$

Bare DM mass is small:
higher order corrections

GWs from bubble collisions

-22-


Bubble wall energy is released to GWs.

i) Bubble collision

ii) Plasma friction

{ sound waves
turbulences

κ_{coll} : efficiency parameters



[K. Schmitz, 2020]

Peak frequency:
$$f_{\text{coll}} = 56.8 \text{ Hz} \left(\frac{\beta/H_*}{10} \right) \left(\frac{T_*}{10^8 \text{ GeV}} \right) \left(\frac{1}{1 - 0.05v_w + 0.55v_w^2} \right) \left(\frac{g_*(T_*)}{100} \right)^{1/6}$$


GW spectrum:
$$\Omega_{\text{coll}}(f) h^2 = 1.66 \times 10^{-5} \left(\frac{H_*}{\beta} \right)^2 \frac{\kappa_{\text{coll}}^2 \alpha^2}{(1 + \alpha)^2} \left(\frac{100}{g_*(T_*)} \right)^{1/3} \frac{v_w^3}{1 + 2.4v_w^2} \cdot \frac{(f/f_{\text{coll}})^{2.8}}{1 + 2.8(f/f_{\text{coll}})^{3.8}}$$

DM vs GW

Experiment	$f_{\text{optimal}}/\text{Hz}$	v_ϕ/GeV	m_{DM}/GeV
Pulsar Timing Arrays (PTAs) [107]	10^{-8}	0.1	$10^{13} - 10^{16}$
LISA [108]	0.001	10^4	$10^6 - 10^{15}$
BBO [109], DECIGO [110]	0.1	10^6	$10^5 - 10^{13}$
Einstein Telescope (ET) [111], Cosmic Explorer (CE) [112]	10	10^8	$10^6 - 10^{10}$

Fermion dark matter

-23-



EFT for fermion DM:

$$\mathcal{L}_{\chi_f} = \frac{1}{\Lambda_f} \phi^2 \bar{\chi}_f \chi_f$$

$$\longrightarrow \mathcal{L}_{\text{eff}} = y_f \phi \bar{\chi}_f \chi_f,$$

$$y_f = \frac{v_\phi}{\Lambda_f}, \quad \Lambda_f < m_{\chi_f}, v_\phi$$

Hot DM, rethermalization,
freeze-in bounds are not
shown: UV-dependent.

$$E_{\chi_f} \gtrsim R_*^3 \Delta V$$

Bare DM mass is small:
higher order corrections

Vector dark matter and gauge choices



Vector dark matter

-24-

Vector DM



$$g^2 \phi^2 V_\mu V^\mu \longrightarrow g^2 (v_\phi + \bar{\phi})^2 V_\mu V^\mu$$

e.g. in unitary gauge



$m_V > g v_\phi \longrightarrow$ Another mechanism for vector DM mass

e.g. mixing with extra Higgs s induces effective interactions.



Gauges for vector DM

-25-

Production of on-shell gauge bosons:

Polarization sum, $\sum \epsilon^\mu \epsilon^\nu \rightarrow -g^{\mu\nu} + (1 - \xi) \frac{p^\mu p^\nu}{p^2 - \xi m_V^2}$

+ ghosts, Goldstone fields with $m^2 = \xi m_V^2$

1) unitary gauge: $\xi \rightarrow \infty$, $\sum \epsilon^\mu \epsilon^\nu \rightarrow -g^{\mu\nu} + \frac{p^\mu p^\nu}{m_V^2}$

$$|\bar{\mathcal{M}}(\phi_p^* \rightarrow VV)|^2 = g^2 m_V^2 \left(3 - \frac{p^2}{m_V^2} + \frac{p^4}{4m_V^4} \right) \quad [\text{A. Falkowski, J.M. No, 2012}]$$

$$p^2 \gg m_\phi^2, m_V^2, \quad |\bar{\mathcal{M}}(\phi_p^* \rightarrow VV)|^2 \sim \frac{g^2}{4m_V^2} p^4$$

2) Feynman-'t Hooft gauge: $\xi = 1$, $\sum \epsilon^\mu \epsilon^\nu \rightarrow -g^{\mu\nu}$

+ ghosts, Goldstone fields with $m_c^2 = m_G^2 = m_V^2$

$$|\bar{\mathcal{M}}(\phi_p^* \rightarrow VV)|^2 = g^2 m_V^2 \left(3 - \frac{p^2}{m_V^2} + \frac{\lambda_\phi^2}{g^4} \right) \longrightarrow \text{still negative } p^2 ?$$

Vector DM in R_ξ gauge

-26-

3) R_ξ gauge: $m_c^2 = m_G^2 = \xi m_V^2 \neq m_V^2$

$$|\bar{\mathcal{M}}(\phi_p^* \rightarrow VV)|^2 = (gm_V)^2 \left(2 + \frac{1}{4m_V^4} (p^2 - 2m_V^2)^2 \right) \sqrt{1 - \frac{4m_V^2}{p^2}}$$

[G. Giudice et al, 2024]


“gauge-independent”

$$+ \frac{g^2}{4m_V^2} (-p^2 + m_\phi^2)(p^2 + m_\phi^2) \sqrt{1 - \frac{4\xi m_V^2}{p^2}}$$

“gauge-dependent”: nonzero of off-shell scalar

$$|\bar{\mathcal{M}}(\phi_p^* \rightarrow VV)|^2 = g^2 m_V^2 \times \begin{cases} \frac{(\xi-3)p^2}{2m_V^2} + \frac{\lambda_\phi^2}{g^4} + 3 & \text{for } \frac{p^2}{m_V^2} \gg \xi, 1 \\ \frac{p^4}{4m_V^4} - \frac{p^2}{m_V^2} + 3 & \text{for } \xi \gg \frac{p^2}{m_V^2}, 1 \end{cases}$$

Fried-Yennie gauge: $\xi = 3 \longrightarrow$ no p^2 or p^4 terms!



off-shell Higgs self-energy: gauge-dependent

$\longrightarrow$ + vertex, box diagrams

cf. [J. Papavasiliou, A. Pilaftsis et al, 1997]


Problem in unitary gauge

-27-

$$p^2 \gg m_\phi^2, m_V^2, \quad |\bar{\mathcal{M}}(\phi_p^* \rightarrow VV)|^2 \sim \frac{g^2}{4m_V^2} p^4 \quad \text{in unitary gauge.}$$

2-body $\phi_p^* \rightarrow 2V$: $\text{Im}[\tilde{\Gamma}^{(2)}(p^2)]_{\phi_p^* \rightarrow 2V} \sim \frac{g^2 m_V^2}{8\pi} \left(\frac{p^4}{4m_V^4} \right)$


4-body $\phi_p^* \rightarrow 4V$:



$$\text{Im}[\tilde{\Gamma}^{(2)}(p^2)]_{\phi_p^* \rightarrow 4V} \sim \frac{g^6 m_V^2}{8\pi(4\pi^2)^2} \left(\frac{p^4}{4m_V^4} \right)^2$$

4-body process “grows faster” with large momentum.

Processes with higher vector boson multiplicity
grows even faster with large momentum.



Breakdown of perturbative expansion!

Goldstone equivalence

-28-

Goldstone Equivalence Theorem:

$$\epsilon_L^\mu \frac{p_\mu}{m_V} = G \quad (\text{Goldstone})$$


Physical processes with longitudinal modes at high energy:

$$\bar{\mathcal{M}}(\phi; V_L) = \bar{\mathcal{M}}(\phi; G) + \mathcal{O}\left(\frac{m_V^2}{p^2}\right) \quad [\text{B.W. Lee et al, 1977}]$$

(i) TT: $|\bar{\mathcal{M}}|^2 \sim 2m_V^2,$

(ii) LL: $\phi_p^* \rightarrow GG, \quad |\bar{\mathcal{M}}|^2 \sim \lambda_\phi^2 v_\phi^2 = \frac{\lambda_\phi^2}{g^2} m_V^2$

(iii) TL: $\phi_p^* \rightarrow V_T(p_1)G(p_2),$



In rest frame for ϕ_p^* , $p = (E, 0, 0, 0)$

V_T, G back-to-back $\longrightarrow |\bar{\mathcal{M}}|^2 \sim 0$


$$|\bar{\mathcal{M}}|^2 \sim m_V^2 \left(2g^2 + \frac{\lambda_\phi^2}{g^2}\right) \left(1 + \mathcal{O}\left(\frac{m_V^2}{p^2}\right)\right), \quad p^2 \gg m_V^2 : \text{finite at large } p^2$$

Other processes

-29-

3-body process for vector DM:

$$\phi_p^* \rightarrow \phi V V$$



VDM part: $|\bar{\mathcal{M}}|_{p^2 < v_\phi^2}^2 \sim v_\phi^2 g^4 \left(3 - \frac{p^2}{m_V^2} + \frac{p^4}{4m_V^4} \right), \quad |\bar{\mathcal{M}}|_{p^2 > v_\phi^2}^2 \sim v_\phi^2 (\lambda_\phi^2 + 2g^4)$

$\longrightarrow \quad \text{Im } \tilde{\Gamma}^{(2)}(p^2) \Big|_{\phi^* \rightarrow \phi V V} \sim p^2 |\bar{\mathcal{M}}|^2 \sim p^2 (\lambda_\phi^2 + 2g^4), \quad p^2 > v_\phi^2$

cf. 2-body process: $\text{Im } \tilde{\Gamma}^{(2)}(p^2) \Big|_{\phi^* \rightarrow V V} \sim |\bar{\mathcal{M}}|^2 \Big|_{\phi^* \rightarrow V V}$

$p^2 > v_\phi^2$: 3-body process dominant for vector DM!

Higher order processes for DM: $\phi_{p_1}^* \phi_{p_2}^* \rightarrow \chi_s \chi_s, \bar{\chi}_f \chi_f, V V$

$\phi_{p_1}^* \sim \left(\frac{v_\phi}{p^2} \right)^2 \longrightarrow \text{Suppressed for } p^2 \sim m_{\text{DM}}^2 \gg v_\phi^2$

Back-reaction issue

-30-

Energy density of particles can be comparable to latent heat:

$$E_{\text{DM}} = 4\pi R_*^2 \int \frac{dp^2}{2\pi^2} p f(p^2) \text{Im} \tilde{\Gamma}^{(2)}(p^2) \sim \frac{4\pi^2}{3} R_*^3 \Delta V$$

→ Bubble dynamics & DM production subject to corrections.

$$f(p^2) \sim \frac{1}{p^4} \quad (\text{Elastic collisions})$$

$$|\bar{\mathcal{M}}|^2 \sim \left\{ \begin{array}{l} (p^2)^0, \text{ (scalar, 2-body)} \quad ; \quad p^2, \text{ (scalar, 3-body)} \\ p^2, \text{ (fermion, 2-body)} \\ \text{--- } p^4, \text{ (Vector, 2-body) --- Back-reaction issue! ---} \\ (p^2)^0, \text{ (vector, 2-body)} \quad ; \quad p^2, \text{ (vector, 3-body)} \end{array} \right.$$

Important to treat the growth of vector
DM properly for back-reaction issue.

Conclusions

- Bubble dynamics in the 1st-order phase transitions is an interesting testing ground for DM and GWs.
- Non-relativistic bubbles: if DM receives mass from SSB,
→ “Filtering mechanism” DM with masses parametrically larger than T .
- Relativistic bubbles: if DM mass is not related to SSB,
→ “Bubble collisions” produce DM with masses much larger than T & v_ϕ .

$$\boxed{m_{\text{DM}} \lesssim \frac{1}{l_w} \sim \gamma_w v_\phi \sim \gamma_w T, \quad \gamma_w \gg 1} \quad \text{cf. wall-plasma: } m_{\text{DM}} \ll \gamma_w T$$

- Vector dark matter is subject to gauge dependence: a practical solution with GET is provided.